

PERMANENCE OF STABLE RANK ONE FOR CENTRALLY LARGE SUBALGEBRAS AND CROSSED PRODUCTS BY MINIMAL HOMEOMORPHISMS

DAWN ARCHEY AND N. CHRISTOPHER PHILLIPS

ABSTRACT. We define centrally large subalgebras of simple unital C^* -algebras, strengthening the definition of large subalgebras in previous work. We prove that if A is any infinite dimensional simple separable unital C^* -algebra which contains a centrally large subalgebra with stable rank one, then A has stable rank one. We also prove that large subalgebras of crossed product type are automatically centrally large. We use these results to prove that if X is a compact metric space which has a surjective continuous map to the Cantor set, and $h: X \rightarrow X$ is a minimal homeomorphism, then $C^*(\mathbb{Z}, X, h)$ has stable rank one, regardless of the dimension of X or the mean dimension of h . In particular, the Giol-Kerr examples give crossed products with stable rank one but which are not stable under tensoring with the Jiang-Su algebra and are therefore not classifiable in terms of the Elliott invariant.

1. INTRODUCTION

Large and stably large subalgebras were introduced in [16] as an abstraction of Putnam's orbit breaking subalgebra of the crossed product $C^*(\mathbb{Z}, X, h)$ of the Cantor set by a minimal homeomorphism. In this paper, we define a stronger concept, that of a centrally large subalgebra. Large orbit breaking subalgebras are also centrally large, and centrally large subalgebras provide more information about the containing algebra. We prove (Theorem 6.3) that if A is a simple unital C^* -algebra and $B \subset A$ is a large subalgebra which has stable rank one, then A also has stable rank one. If in addition B has real rank zero, we prove (Theorem 6.4) that A also has real rank zero.

We use our result to prove the following result (Theorem 7.1) for crossed products by minimal homeomorphisms. Let X be a compact metric space which has a surjective continuous map to the Cantor set, and let $h: X \rightarrow X$ be a minimal homeomorphism. Then $C^*(\mathbb{Z}, X, h)$ has stable rank one. This result holds regardless of the dimension of X or the mean dimension of h . It is expected that $C^*(\mathbb{Z}, X, h)$ is classifiable in terms of the Elliott invariant if and only if h has mean dimension zero, and one direction is the main result of [9]. Thus our result shows that crossed products by minimal homeomorphisms can have stable rank one even when they are not expected to be classifiable. The examples of [10] confirm that this happens: they do not have mean dimension zero, their crossed products are

Date: 30 March 2016.

2010 *Mathematics Subject Classification.* Primary 46L05; Secondary 46L55.

This material is partially based upon work of the second author supported by the US National Science Foundation under Grants DMS-0701076 and DMS-1101742. It was also partially supported by the Centre de Recerca Matemàtica (Barcelona) through research visits by both authors conducted during 2011.

not stable under tensoring with the Jiang-Su algebra, but our theorem shows that the crossed products do have stable rank one.

In [18], we will use our result to prove a generalization of Theorem 7.1. Let X be a compact metric space with a free minimal action of $\mathbb{Z}^d$. Assume $(X, \mathbb{Z}^d)$ has a factor system which is a free minimal action of $\mathbb{Z}^d$ on the Cantor set. Then $C^*(\mathbb{Z}, X)$ has stable rank one. Again, no hypothesis on mean dimension is needed. We do not include the proof here because the proof of the existence of the necessary centrally large subalgebra is somewhat involved. We point out, however, that the main reason for the abstraction of Putnam's construction is to be able to handle situations of this sort, in which there appears to be no explicit construction of the necessary subalgebra, only an existence proof.

We give the motivation and background in more detail. Let X be an infinite compact metric space, let $h: X \rightarrow X$ be a homeomorphism, and let $Y \subset X$ be closed. Let $u \in C^*(\mathbb{Z}, X, h)$ be the standard unitary. The Y -orbit breaking subalgebra is the subalgebra $C^*(\mathbb{Z}, X, h)_Y$ of $C^*(\mathbb{Z}, X, h)$ generated by $C(X)$ and all elements fu for $f \in C(X)$ such that $f|_Y = 0$. (Putnam actually used uf rather than fu , but our choice makes the relationship with Rokhlin towers less awkward.) If h is minimal, X is infinite, and Y meets each orbit of h at most once, then $C^*(\mathbb{Z}, X, h)_Y$ is large in $C^*(\mathbb{Z}, X, h)$ (Theorem 7.10 of [16]).

Subalgebras with properties similar to this one have been used in many other places, such as the study of crossed products of the Cantor set by free minimal actions of $\mathbb{Z}^d$ ([15], but without the name), the recent proof by Elliott and Niu that if $h: X \rightarrow X$ is minimal and has mean dimension zero, then $C^*(\mathbb{Z}, X, h)$ is Z -stable [9], and the proof that “breaking” the generating unitaries in an irrational rotation algebra gives an AF algebra [8]. See the introduction to [16] for a much longer list of applications.

For an infinite dimensional simple unital C^* -algebra A and a stably large subalgebra $B \subset A$, the following results were proved in [16]:

- B is simple (Proposition 5.2 of [16]) and infinite dimensional (Proposition 5.5 of [16]).
- If B is stably finite then so is A (Corollary 6.16 of [16]), and if B is purely infinite then so is A (Proposition 6.17 of [16]).
- The restriction maps $T(A) \rightarrow T(B)$ and $QT(A) \rightarrow QT(B)$ (on tracial states and quasitraces) are bijective (Theorem 6.2 of [16] and Proposition 6.9 of [16]).
- When A is stably finite, the inclusion of B in A induces an isomorphism on the semigroups that remain after deleting from $Cu(B)$ and $Cu(A)$ all the classes of nonzero projections (Theorem 6.8 of [16]).
- B and A have the same radius of comparison (Theorem 6.14 of [16]).

However, for several further results, such as ones involving stable rank one and Z -stability, one seems to need a stronger condition: B must be what we call centrally large in A .

In this paper, we define centrally large subalgebras. We prove that a large subalgebra of crossed product type (Definition 4.1 below; Definition 4.9 of [16]) is in fact centrally large; this result covers most of the examples which have appeared in applications. We then prove that if B is centrally large in A and B has stable rank one, then so does A . We will consider Z -stability elsewhere [1].

For the application to the stable rank of crossed products, let $h: X \rightarrow X$ be as above, and assume that X has a surjective continuous map to the Cantor set. (The Giol-Kerr examples [10] have this property, because they have the universal odometer as a factor.) Then Y as above can be chosen to be a decreasing intersection of compact open subsets of X . In this case, it is known that $C^*(\mathbb{Z}, X, h)_Y$ is the direct limit of an AH system which has diagonal maps in the sense of [7]. Moreover (by [7]), a simple direct limit of an AH system with diagonal maps has stable rank one, even with no assumptions on the dimension growth in the system.

We conjecture, jointly with Zhuang Niu, that $C^*(\mathbb{Z}, X, h)$ has stable rank one for a completely arbitrary minimal homeomorphism h of an infinite compact metric space X .

The original main application was to prove that if X is a finite dimensional compact metric space, and if $\mathbb{Z}^d$ acts freely and minimally on X , then $C^*(\mathbb{Z}^d, X)$ has stable rank one, using the large subalgebra of crossed product type constructed in [17]. This particular application has been superseded before its publication in two different ways. First, using centrally large subalgebras, one can in fact prove Z -stability, which by Theorem 6.7 of [25] implies stable rank one. Second, Z -stability (in fact, finite nuclear dimension) has also been obtained by different methods in [27].

The paper is organized as follows. In Section 2 we establish notation and prove some preliminary lemmas on approximation and Cuntz comparison. In Section 3 we give the definition of a centrally large subalgebra and some variations on the theme. In Section 4 we prove that large subalgebras of crossed product type are centrally large. In Section 5 we prove some technical lemmas. In Section 6 we prove the main theorem and the result about real rank zero. Section 7 contains the result on the stable rank of $C^*(\mathbb{Z}, X, h)$ and its application to examples, including those of Giol and Kerr.

We use the following notational conventions. If A is a C^* -algebra, then A_+ denotes the set of positive elements of A . Also, A^+ denotes the unitization of A . (We add a new identity even if A is already unital.) The notations $M_\infty(A)$ and $a \oplus b$ for $a, b \in M_\infty(A)$ and $a, b \in K \otimes A$ are described in Section 2. By convention, if we say that B is a unital subalgebra of a C^* -algebra A , we mean that B contains the identity of A . Subalgebras of C^* -algebras are always supposed to be C^* -subalgebras. We take $[a, b] = ab - ba$.

We are grateful to Francesc Perera for pointing out the reference [26], to Zhuang Niu for pointing out the reference [7], and to George Elliott for pointing out Corollary 5 of [3].

2. CUNTZ COMPARISON AND APPROXIMATION LEMMAS

We give definitions and notation related to Cuntz comparison, and one result needed later for which we don't know a reference. We refer to the list in Lemma 1.4 of [16], and the results elsewhere in Section 1 of [16], for many more facts. Also see the survey article [2]. At the end, we prove several approximation lemmas which will be needed later.

For a C^* -algebra A , let $M_\infty(A)$ denote the algebraic direct limit of the system $(M_n(A))_{n=1}^\infty$ using the usual embeddings $M_n(A) \rightarrow M_{n+1}(A)$, given by

$$a \mapsto \begin{pmatrix} a & 0 \\ 0 & 0 \end{pmatrix}.$$

If $a \in M_m(A)$ and $b \in M_n(A)$, we write $a \oplus b$ for the diagonal direct sum

$$a \oplus b = \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}.$$

By abuse of notation, we will also write $a \oplus b$ when $a, b \in M_\infty(A)$ and we do not care about the precise choice of m and n with $a \in M_m(A)$ and $b \in M_n(A)$. We further choose some isomorphism $M_2(K) \rightarrow K$, and for $a, b \in K \otimes A$ we use the resulting isomorphism $M_2(K \otimes A) \rightarrow K \otimes A$ to interpret $a \oplus b$ as an element of $K \otimes A$. Up to unitary equivalence which is trivial on A , the result does not depend on the choice of the isomorphism $M_2(K) \rightarrow K$.

The following definitions are originally from [5].

Definition 2.1. Let A be a C^* -algebra, and let $a, b \in (K \otimes A)_+$.

- (1) We say that a is Cuntz subequivalent to b over A , written $a \preceq_A b$, if there is a sequence $(v_n)_{n=1}^\infty$ in $K \otimes A$ such that

$$\lim_{n \rightarrow \infty} v_n b v_n^* = a.$$

- (2) We say that a and b are Cuntz equivalent in A , written $a \sim_A b$, if $a \preceq_A b$ and $b \preceq_A a$.

The relation $\sim_A$ is an equivalence relation. When there is no possible confusion about the algebra A , we suppress it in the notation.

We will regularly use the following two families of functions.

Definition 2.2. Let A be a C^* -algebra, let $a \in A_+$, and let $\varepsilon > 0$. We define $(a - \varepsilon)_+$ as follows. Let $f: [0, \infty) \rightarrow [0, \infty)$ be the function

$$f(\lambda) = (\lambda - \varepsilon)_+ = \begin{cases} 0 & 0 \leq \lambda \leq \varepsilon \\ \lambda - \varepsilon & \varepsilon < \lambda. \end{cases}$$

Then define $(a - \varepsilon)_+ = f(a)$ (using continuous functional calculus).

Definition 2.3. For $\varepsilon > 0$, define a continuous function $f_\varepsilon: [0, \infty) \rightarrow [0, 1]$ by

$$f_\varepsilon(\lambda) = \begin{cases} 0 & 0 \leq \lambda \leq \varepsilon \\ \varepsilon^{-1}(\lambda - \varepsilon) & \varepsilon \leq \lambda \leq 2\varepsilon \\ 1 & 2\varepsilon \leq \lambda. \end{cases}$$

Lemma 2.4. Let A be a simple unital C^* -algebra, and let $a, b \in A_+$ with $\|a\| = \|b\| = 1$. Then there exists $c \in A_+ \setminus \{0\}$ with $\|c\| \leq 1$ such that $c \leq a$ and $c \preceq b$.

Proof. Since A is simple and $a, b \in A$ are nonzero, by Proposition 1.8 of [4] there is a nonzero $y \in A$ such that $yy^* \in \overline{aAa}$ and $y^*y \in \overline{bAb}$. Without loss of generality we may assume that $\|y\| \leq 1$, and so $yy^* \leq 1$. Set $z = y^*a^{1/2}$ and $c = z^*z$. Then $c = a^{1/2}yy^*a^{1/2} \leq a$. Also, the discussion after Definition 2.3 of [12] gives the first step in the calculation $c \sim zz^* = y^*ay \leq yy^* \in \overline{bAb}$, so $c \preceq b$ by Proposition 2.7(i) of [12]. ■

We finish this section with several approximation lemmas. The first is actually a special case of Lemma 2.5.11(1) of [13], but the proof there is a bit sloppy.

Lemma 2.5. *Suppose $f: [0, 1] \rightarrow \mathbb{C}$ is continuous. Then for every $\varepsilon > 0$ there exists $\delta > 0$ such that whenever D is a C^* -algebra and $x, z \in D$ satisfy*

$$0 \leq x \leq 1, \quad \|z\| \leq 1, \quad \text{and} \quad \|[x, z]\| < \delta,$$

then $\|[f(x), z]\| < \varepsilon$.

Proof. We first observe that if $\|x\| \leq 1$ and $\|z\| \leq 1$, then for $n \in \mathbb{Z}_{>0}$ we have

$$(2.1) \quad \|x^n, z\| \leq \sum_{k=1}^n \|x^{n-k}\| \cdot \|[x, z]\| \cdot \|x^{k-1}\| \leq n\|[x, z]\|.$$

Choose $n \in \mathbb{Z}_{\geq 0}$ and a polynomial $g(\lambda) = \sum_{k=0}^n \alpha_k \lambda^k$ with coefficients $\alpha_k \in \mathbb{C}$ for $k = 0, 1, \dots, n$ such that $|g(\lambda) - f(\lambda)| < \frac{\varepsilon}{3}$ for all $\lambda \in [0, 1]$. Define

$$\delta = \frac{\varepsilon}{1 + 3 \sum_{k=1}^n k |\alpha_k|}.$$

Suppose $\|[x, z]\| < \delta$. Then, since $\|z\| \leq 1$ and using (2.1) at the second step,

$$\|[f(x), z]\| \leq 2\|f(x) - g(x)\| + \sum_{k=1}^n |\alpha_k| \cdot \|[x^k, z]\| < 2 \left(\frac{\varepsilon}{3} \right) + \sum_{k=1}^n k |\alpha_k| \delta < \varepsilon.$$

This completes the proof. $\blacksquare$

Lemma 2.6. *Suppose $f: [0, 1] \rightarrow \mathbb{C}$ is continuous and $f(0) = 0$. Then for every $\varepsilon > 0$ there exists $\delta > 0$ such that whenever D is a C^* -algebra, $E \subset D$ is a subalgebra, and $a \in D$ and $b \in E$ satisfy*

$$\|a\| \leq 1, \quad 0 \leq b \leq 1, \quad \text{and} \quad \text{dist}(ba, E) < \delta,$$

then $\text{dist}(f(b)a, E) < \varepsilon$.

Proof. Choose $n \in \mathbb{Z}_{\geq 0}$ and a polynomial $g(\lambda) = \sum_{k=0}^n \alpha_k \lambda^k$ with $\alpha_k \in \mathbb{C}$ for $k = 0, 1, \dots, n$ such that $\alpha_0 = 0$ and $|g(\lambda) - f(\lambda)| < \frac{\varepsilon}{3}$ for all $\lambda \in [0, 1]$. Define

$$\delta = \frac{\varepsilon}{2(1 + |\alpha_1| + |\alpha_2| + \dots + |\alpha_n|)}.$$

Let $a \in D$ and $b \in E$ be as in the hypotheses. Choose $y \in E$ such that $\|ba - y\| < \delta$. Set $z = \sum_{k=1}^n \alpha_k b^{k-1} y \in E$. Then

$$\|f(b)a - z\| \leq \|f(b) - g(b)\| \cdot \|a\| + \sum_{k=1}^n |\alpha_k| \cdot \|b^{k-1}\| \cdot \|ba - y\| < \frac{\varepsilon}{2} + \sum_{k=1}^n |\alpha_k| \delta < \varepsilon.$$

This completes the proof. $\blacksquare$

The following corollary is also easy to prove directly.

Corollary 2.7. *Let D be a C^* -algebra, let $E \subset D$ be a subalgebra, let $a \in D$, let $b \in E_+$, and suppose that $ba \in E$. Let $f: \text{sp}(b) \rightarrow \mathbb{C}$ be a continuous function such that $f(0) = 0$. Then $f(b)a \in E$.*

Proof. By scaling, without loss of generality $\|b\| \leq 1$. Then we may extend f to be a continuous function $f: [0, 1] \rightarrow \mathbb{C}$. Lemma 2.6 now shows that for all $\varepsilon > 0$, we have $\text{dist}(f(b)a, E) < \varepsilon$. $\blacksquare$

3. DEFINITIONS AND PRELIMINARY LEMMAS

In this section, we recall the definition of a large subalgebra, and give the definition of a centrally large subalgebra. We then give some convenient variants of the definition.

Recall that a unital subalgebra of a unital C^* -algebra A is, by convention, assumed to contain the identity of A .

The definitions of “large” and “stably large” in Definition 3.1 originally appeared in Definition 4.1 and Definition 5.1 of [16]. The definitions of “centrally large” and “stably centrally large” differ by the addition of an approximate commutation condition.

Definition 3.1. *Let A be an infinite dimensional simple unital C^* -algebra. A unital subalgebra $B \subset A$ is said to be large in A if for every $m \in \mathbb{Z}_{>0}$, $a_1, a_2, \dots, a_m \in A$, $\varepsilon > 0$, $x \in A_+$ with $\|x\| = 1$, and $y \in B_+ \setminus \{0\}$, there are $c_1, c_2, \dots, c_m \in A$ and $g \in B$ such that:*

- (1) $0 \leq g \leq 1$.
- (2) For $j = 1, 2, \dots, m$ we have $\|c_j - a_j\| < \varepsilon$.
- (3) For $j = 1, 2, \dots, m$ we have $(1 - g)c_j \in B$.
- (4) $g \preceq_B y$ and $g \preceq_A x$.
- (5) $\|(1 - g)x(1 - g)\| > 1 - \varepsilon$.

We say that B is centrally large in A if we can require that in addition:

- (6) For $j = 1, 2, \dots, m$ we have $\|ga_j - a_jg\| < \varepsilon$.

We further say that B is stably large in A if $M_n(B)$ is large in $M_n(A)$ for all $n \in \mathbb{Z}_{\geq 0}$, and stably centrally large in A if $M_n(B)$ is centrally large in $M_n(A)$ for all $n \in \mathbb{Z}_{\geq 0}$.

We emphasize that one of the Cuntz subequivalences in (4) is relative to B , not to A .

By Proposition 4.5 of [16], when A is finite we can omit the condition involving $\|(1 - g)x(1 - g)\|$ in the definition of a large subalgebra. The following proposition gives the same result for centrally large subalgebras.

Proposition 3.2. *Let A be an infinite dimensional simple unital C^* -algebra, and let $B \subset A$ be a unital subalgebra. Assume that A is finite and that for every $m \in \mathbb{Z}_{>0}$, $a_1, a_2, \dots, a_m \in A$, $\varepsilon > 0$, and $y \in B_+ \setminus \{0\}$, there are $c_1, c_2, \dots, c_m \in A$ and $g \in B$ such that:*

- (1) $0 \leq g \leq 1$.
- (2) For $j = 1, 2, \dots, m$ we have $\|c_j - a_j\| < \varepsilon$.
- (3) For $j = 1, 2, \dots, m$ we have $(1 - g)c_j \in B$.
- (4) $g \preceq_B y$ and $g \preceq_A x$.
- (5) For $j = 1, 2, \dots, m$ we have $\|ga_j - a_jg\| < \varepsilon$.

Then B is a centrally large subalgebra of A .

Proof. The proof is the same as that of Proposition 4.5 of [16]. ■

The following lemma states that the elements c_j in the definition of a centrally large subalgebra can be chosen with norm no larger than the norm of corresponding a_j .

Lemma 3.3. *Suppose $B \subset A$ is a centrally large subalgebra of A . The elements $c_1, c_2, \dots, c_m$ may be chosen so that $\|c_j\| \leq \|a_j\|$ for $j = 1, 2, \dots, m$.*

Proof. The proof is the same as the proof of Lemma 4.7 in [16]. ■

If we cut down on both sides instead of on one side, and the elements a_j are positive, then we may take the elements c_j to be positive. We can still choose c_j with norm no larger than that of a_j .

Lemma 3.4. *Let A be an infinite dimensional simple unital C^* -algebra. Suppose $B \subset A$ is a centrally large subalgebra. Let $m, n \in \mathbb{Z}_{\geq 0}$, let $a_1, a_2, \dots, a_m \in A$, let $b_1, b_2, \dots, b_n \in A_+$, let $\varepsilon > 0$, let $x \in A_+$ satisfy $\|x\| = 1$, and let $y \in B_+ \setminus \{0\}$. Then there are $c_1, c_2, \dots, c_m \in A$, $d_1, d_2, \dots, d_n \in A_+$, and $g \in B$ such that:*

- (1) $0 \leq g \leq 1$.
- (2) For $j = 1, 2, \dots, m$ we have $\|c_j - a_j\| < \varepsilon$, and for $j = 1, 2, \dots, n$ we have $\|d_j - b_j\| < \varepsilon$.
- (3) For $j = 1, 2, \dots, m$ we have $\|c_j\| \leq \|a_j\|$, and for $j = 1, 2, \dots, n$ we have $\|d_j\| \leq \|b_j\|$.
- (4) For $j = 1, 2, \dots, m$ we have $(1 - g)c_j \in B$, and for $j = 1, 2, \dots, n$ we have $(1 - g)d_j(1 - g) \in B$.
- (5) $g \preceq_B y$ and $g \preceq_A x$.
- (6) $\|(1 - g)x(1 - g)\| > 1 - \varepsilon$.
- (7) $\|ga_j - a_jg\| < \varepsilon$ for $j = 1, 2, \dots, m$ and $\|gb_j - b_jg\| < \varepsilon$ for $j = 1, 2, \dots, n$.

Proof. The proof is the essentially the same as the proof of Lemma 4.8 in [16]. ■

Proposition 3.5. *Let A_1 and A_2 be infinite dimensional simple unital C^* -algebras, and let $B_1 \subset A_1$ and $B_2 \subset A_2$ be centrally large subalgebras. Assume that $A_1 \otimes_{\min} A_2$ is finite. Then $B_1 \otimes_{\min} B_2$ is a centrally large subalgebra of $A_1 \otimes_{\min} A_2$.*

Proof. The proof is essentially the same as that of Proposition 5.6 of [16]; we describe the differences.

In place of Proposition 4.5 of [16], we use Proposition 3.2. The analog for centrally large subalgebras of Lemma 4.2 of [16] is obvious.

We also need to check the approximate commutation relation in the conclusion, that is, using the notation of the proof of Proposition 5.6 of [16], we take

$$g = 1 - (1 - g_1) \otimes (1 - g_2),$$

and we need to prove that

$$\|[g, a_{1,j} \otimes a_{2,j}]\| < \varepsilon$$

for $j = 1, 2, \dots, m$. For $l = 1, 2$, since B_l is centrally large in A_l , we may require, in addition to the conditions demanded in the proof of Proposition 5.6 of [16], that $\|[g_l, a_{l,j}]\| < \frac{\varepsilon}{2}$. Then also $\|[1 - g_l, a_{l,j}]\| < \frac{\varepsilon}{2}$. Therefore, using $\|1 - g_l\| \leq 1$ and $\|a_{l,j}\| \leq 1$ at the fourth step, we get

$$\begin{aligned} \|[g, a_{1,j} \otimes a_{2,j}]\| &= \|[1 - g, a_{1,j} \otimes a_{2,j}]\| = \|[(1 - g_1) \otimes (1 - g_2), a_{1,j} \otimes a_{2,j}]\| \\ &\leq \|[1 - g_1, a_{1,j}]\| \otimes \|(1 - g_2)a_{2,j}\| + \|a_{1,j}(1 - g_1) \otimes [1 - g_2, a_{2,j}]\| \\ &< \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon. \end{aligned}$$

This completes the proof. ■

We now get the analog of Corollary 5.8 of [16].

Corollary 3.6. *Let A be a stably finite infinite dimensional simple unital C^* -algebra, and let $B \subset A$ be a centrally large subalgebra. Then B is stably centrally large in A .*

Proof. In Proposition 3.5 take $A_1 = B_1 = M_n$, $A_2 = A$, and $B_2 = B$. ■

In the definition of a centrally large subalgebra, it suffices to verify the conditions for $a_1, a_2, \dots, a_m$ in a generating set. (We do not know the analog of this statement for large subalgebras. The approximate commutation relation makes the difference.) We do not use this result in the rest of the paper, but it seems potentially useful elsewhere.

The proof requires the analog of Proposition 4.4 of [16].

Lemma 3.7. *Let A be an infinite dimensional simple unital C^* -algebra, and let $B \subset A$ be a unital subalgebra. Suppose that every finite set $F \subset A$, $\varepsilon > 0$, $x \in A_+$ with $\|x\| = 1$, and $y \in B_+ \setminus \{0\}$, there is $g \in B$ such that:*

- (1) $0 \leq g \leq 1$.
- (2) $\text{dist}((1-g)a, B) < \varepsilon$ for all $a \in F$.
- (3) $g \preceq_B y$ and $g \preceq_A x$.
- (4) $\|(1-g)x(1-g)\| > 1 - \varepsilon$.
- (5) $\|ga - ag\| < \varepsilon$ for all $a \in F$.

Then B is centrally large in A .

Proof. The proof is essentially the same as that of Proposition 4.4 of [16], except with one addition. We describe only the addition.

We verify the conditions of Definition 3.1. Let $m \in \mathbb{Z}_{\geq 0}$, let $a_1, a_2, \dots, a_m \in A$, let $\varepsilon > 0$, let $x \in A_+$ satisfy $\|x\| = 1$, and let $y \in B_+ \setminus \{0\}$. Let the functions $f_0, f_1, f_2: [0, 1] \rightarrow [0, 1]$ be as in the proof of Proposition 4.4 of [16]. Define a continuous function $f: [0, 1] \rightarrow [0, 1]$ by $f(\lambda) = 1 - f_0(1 - \lambda)$ for $\lambda \in [0, 1]$. Apply Lemma 2.5 with this function f and with ε as given, getting $\delta > 0$. Apply the hypothesis with $F = \{a_1, a_2, \dots, a_m\}$ and with $\min(\delta, \frac{\varepsilon}{3})$ in place of ε , getting $g_0 \in B$. (The difference is that in [16] we used $\frac{\varepsilon}{3}$ here.) Define $r_0 = 1 - g_0$. Set $g = 1 - f_2(r_0)$.

For $j = 1, 2, \dots, m$, define c_j as in the proof of Proposition 4.4 of [16]. The verification of (1), (2), (3), (4), and (5) of Definition 3.1 is exactly as in [16]. To verify Definition 3.1(6), for $j = 1, 2, \dots, m$ we use $\|g_0 a_j - a_j g_0\| < \delta$, $\|a_j\| \leq 1$, and the choice of δ , to get $\|g a_j - a_j g\| < \varepsilon$. ■

Proposition 3.8. *Let A be an infinite dimensional simple unital C^* -algebra, and let $B \subset A$ be a unital subalgebra. Let $S \subset A$ be a subset which generates A as a C^* -algebra. Suppose that every finite set $F \subset S$, $\varepsilon > 0$, $x \in A_+$ with $\|x\| = 1$, and $y \in B_+ \setminus \{0\}$, there is $g \in B$ such that the conditions (1), (2), (3), (4), and (5) of Lemma 3.7 hold. Then B is centrally large in A .*

Proof. It is clear that if $T \subset A$ spans a dense subset of A , then Lemma 3.7 is still valid if in the hypotheses we restrict to finite sets $F \subset T$.

Now suppose that $S \subset A$ generates A as a norm closed subalgebra (without using the adjoint), and the hypotheses of the proposition hold for this set S . We prove that B is centrally large in A by verifying the hypotheses of the generalization of Lemma 3.7 as in the first paragraph. Without loss of generality $\|a\| \leq 1$ for all

$a \in S$. If the hypotheses hold for S , they also hold for $S \cup \{1\}$, so we may assume $1 \in S$.

Let T be the set of all finite products of elements of S . Following the first paragraph, let $F \subset T$ be finite, let $\varepsilon > 0$, let $x \in A_+$ satisfy $\|x\| = 1$, and let $y \in B_+ \setminus \{0\}$; it is enough to show that there is $g \in B$ such that the conditions (1), (2), (3), (4), and (5) of Lemma 3.7 hold. Without loss of generality $\varepsilon < 1$.

Since $1 \in S$, there are $n \in \mathbb{Z}_{>0}$ and a finite set $E \subset S$ such that F is contained in the finite set

$$F_0 = \{a_1 a_2 \cdots a_n : a_1, a_2, \dots, a_n \in E\}.$$

We verify the conditions of Lemma 3.7 for F_0 instead of F .

Use Lemma 2.5 to choose $\delta_1 > 0$ such that whenever $g \in A_+$ and $a \in A$ satisfy

$$0 \leq g \leq 1, \quad \|a\| \leq 1, \quad \text{and} \quad \|ga - ag\| < \delta_1,$$

then

$$(3.1) \quad \|(1-g)^{1/n}a - a(1-g)^{1/n}\| < \frac{\varepsilon}{n^2}.$$

Use Lemma 2.6 (taking $b = 1 - g$) to choose $\delta_2 > 0$ such that whenever $g \in B_+$ and $a \in A$ satisfy

$$\|a\| \leq 1, \quad 0 \leq g \leq 1, \quad \text{and} \quad \text{dist}((1-g)a, B) < \delta_2,$$

then

$$\text{dist}((1-g)^{1/n}a, B) < \frac{\varepsilon}{2^n n}.$$

Set $\delta = \min(\delta_1, \delta_2, \frac{\varepsilon}{n})$. Apply the assumption with E in place of F , with δ in place of ε , and with x and y as given, obtaining $g \in B$.

Conditions (1), (3), and (4) are immediate. For (5), let $a_1, a_2, \dots, a_n \in E$. We use $\|ga_j - a_jg\| < \frac{\varepsilon}{n}$ and $\|a_j\| \leq 1$ for $j = 1, 2, \dots, n$ to get

$$(3.2) \quad \|[g, a_1 a_2 \cdots a_n]\| \leq \sum_{j=1}^n \|a_1 a_2 \cdots a_{j-1}\| \cdot \|ga_j - a_jg\| \cdot \|a_{j+1} a_{j+2} \cdots a_n\| < \varepsilon.$$

We verify (2). Using (3.1), an estimate similar to that for (3.2) gives

$$\|[(1-g)^{1/n}, a_1 a_2 \cdots a_k] a_{k+1} a_{k+2} \cdots a_n\| < \frac{k\varepsilon}{n^2}$$

for $k = 1, 2, \dots, n$. Using $((1-g)^{1/n})^n = 1 - g$ and

$$\|1 - g\|, \|a_1\|, \|a_2\|, \dots, \|a_n\| \leq 1,$$

we then get

$$(3.3) \quad \|(1-g)a_1 a_2 \cdots a_n - (1-g)^{1/n} a_2 (1-g)^{1/n} a_1 \cdots (1-g)^{1/n} a_n\| < \sum_{k=0}^{n-1} \frac{k\varepsilon}{n^2} < \frac{\varepsilon}{2}.$$

Now for $j = 1, 2, \dots, n$, use $\text{dist}((1-g)a_j, B) < \delta_2$ and the choice of δ_2 to choose $b_j \in B$ such that

$$\|(1-g)^{1/n}a_j - b_j\| < \frac{\varepsilon}{2^n n}.$$

Since $\varepsilon < 1$ and $\|a_j\| \leq 1$, we have $\|b_j\| \leq 2$; also, $\|(1-g)^{1/n}a_j\| \leq 1 \leq 2$. For $k = 1, 2, \dots, n$, we therefore get

$$\begin{aligned} & \|b_1 b_2 \cdots b_{k-1} [(1-g)^{1/n}a_k - b_k] (1-g)^{1/n}a_{k+1} (1-g)^{1/n}a_{k+2} \cdots (1-g)^{1/n}a_n\| \\ & < 2^{n-1} \left(\frac{\varepsilon}{2^n n} \right) = \frac{\varepsilon}{2n}. \end{aligned}$$

It follows that

$$\|b_1 b_2 \cdots b_n - (1-g)^{1/n}a_1 (1-g)^{1/n}a_2 (1-g)^{1/n}a_n\| < \frac{\varepsilon}{2}.$$

Since $b_1 b_2 \cdots b_n \in B$, combining this estimate with (3.3) gives

$$\text{dist}((1-g)a_1 a_2 \cdots a_n, B) < \varepsilon.$$

This completes the proof of (2), and the verification of the proposition under the assumption that S generates A as a norm closed subalgebra.

We now prove the proposition as stated by reducing the general case to the case just done. Without loss of generality $\|a\| \leq 1$ for all $a \in S$. Set $T = S \cup S^*$. Let $\varepsilon > 0$, let $F \subset T$ be finite, let $\varepsilon > 0$, let $x \in A_+$ satisfy $\|x\| = 1$, and let $y \in B_+ \setminus \{0\}$. Choose g as in the hypotheses with $\frac{\varepsilon}{2}$ in place of ε . Conditions (1), (3), and (4) are immediate. Conditions (2) and (5) for elements of S are immediate. So let $a \in S^*$. Thus $a^* \in S$, so $g^* = g$ implies

$$\|ga - ag\| = \|a^*g - ga^*\| < \frac{\varepsilon}{2} < \varepsilon,$$

which is (5). Also,

$$\text{dist}(ag, B) = \text{dist}((ag)^*, B) = \text{dist}(ga^*, B) < \frac{\varepsilon}{2},$$

so

$$\text{dist}(ga, B) \leq \|ga - ag\| + \text{dist}(ag, B) < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon,$$

which is (2). ■

4. LARGE SUBALGEBRAS OF CROSSED PRODUCT TYPE

The motivating example for centrally large subalgebras (Definition 3.1) is subalgebras which arise in the study of crossed products by minimal homeomorphisms. In particular, examples of centrally large subalgebras are provided by orbit breaking subalgebras $C^*(\mathbb{Z}, X, h)_Y$ as in the introduction, under the condition that $h^n(Y) \cap Y = \emptyset$ for all $n \in \mathbb{Z} \setminus \{0\}$. These are generalizations of Putnam's algebra $C^*(\mathbb{Z}, X, h)_{\{y\}} \subset C^*(\mathbb{Z}, X, h)$ from [20], and the condition above, that Y meet each orbit at most once, was used by Putnam in Example 2.6 of [21]. See the discussion after Definition 7.3 in [16] for more, and for further generalizations. These are in fact large subalgebras of crossed product type (Definition 4.1 below). Definition 4.1 has the advantage of not explicitly requiring any commutation relations.

Definition 4.1 (Definition 4.9 of [16]). *Let A be an infinite dimensional simple separable unital C^* -algebra. A subalgebra $B \subset A$ is said to be a large subalgebra of crossed product type if there exist a subalgebra $C \subset B$ and a subset G of the unitary group of A such that:*

- (1) (a) C contains the identity of A .
- (b) C and G generate A as a C^* -algebra.
- (c) $uC u^* \subset C$ and $u^* C u \subset C$ for all $u \in G$.

- (2) For every $m \in \mathbb{Z}_{>0}$, $a_1, a_2, \dots, a_m \in A$, $\varepsilon > 0$, $x \in A_+$ with $\|x\| = 1$, and $y \in B_+ \setminus \{0\}$, there are $c_1, c_2, \dots, c_m \in A$ and $g \in C$ such that:
- (a) $0 \leq g \leq 1$.
 - (b) For $j = 1, 2, \dots, m$ we have $\|c_j - a_j\| < \varepsilon$.
 - (c) For $j = 1, 2, \dots, m$ we have $(1 - g)c_j \in B$.
 - (d) $g \preceq_B y$ and $g \preceq_A x$.
 - (e) $\|(1 - g)x(1 - g)\| > 1 - \varepsilon$.

The conditions in (2) are the same as the conditions in Definition 3.1; the difference is that we require that $g \in C$, not merely that $g \in B$. In particular, we have the following relation between the properties.

Proposition 4.2 (Proposition 4.10 of [16]). *Let A be an infinite dimensional simple separable unital C^* -algebra, and let $B \subset A$ be a subalgebra. If B is a large subalgebra of A of crossed product type in the sense of Definition 4.1, then B is a large subalgebra of A in the sense of Definition 3.1.*

Proof. This is immediate from the definitions. ■

In the basic examples, G will be a discrete amenable group, A will have the form $A = C^*(G, X)$ for some essentially free minimal action of G on a compact metric space X , the set G in the definition will be the standard copy of the group G in $C^*(G, X)$, the subalgebra C will be $C(X)$, and B will be a subalgebra of A which contains $C(X)$ but is, informally, much closer to A than to $C(X)$. However, in the definition we do not require that C be commutative, and we do not require that G be a group. In particular, the definition may be applicable to crossed products of noncommutative C^* -algebras, to cocycle crossed products, and to quotients of some nonsimple crossed products. However, something more complicated may well be needed for use with groupoid C^* -algebras or more general crossed products.

The main result of this section is that large subalgebras of crossed product type are centrally large. To improve readability, we isolate several parts of the argument as separate lemmas.

Lemma 4.3. *Let A be a C^* -algebra, let $\varepsilon, \rho \geq 0$, and let $a, b, d, r, x, y \in A_{sa}$ satisfy the following:*

$$a \leq d \leq b, \quad \|b - a\| \leq \varepsilon, \quad \|a - x\| \leq \rho, \quad \|r - y\| \leq \rho, \quad x \leq r, \quad \text{and} \quad y \leq b.$$

Then $\|d - r\| \leq \varepsilon + \rho$.

Proof. Subtracting d from all the elements listed, we reduce to the case $d = 0$, and we must prove that $\|r\| \leq \varepsilon + \rho$.

Since $\|a - x\| \leq \rho$, we have $a - \rho \leq x$. From $\|a - b\| \leq \varepsilon$ and $b \geq 0$, we get $-\varepsilon \leq a - b \leq a$. So $-\varepsilon - \rho \leq a - \rho \leq x \leq r$.

From $\|r - y\| \leq \rho$ we get $r \leq y + \rho$. Since $a \leq 0$ and $\|b - a\| \leq \varepsilon$, we have $y \leq b \leq b - a \leq \varepsilon$. Therefore $r \leq \varepsilon + \rho$. Combining this with the conclusion of the previous paragraph, we get $\|r\| \leq \varepsilon + \rho$, as desired. ■

Lemma 4.4. *For every $\varepsilon > 0$ there is $\delta > 0$ such that whenever A is a C^* -algebra and $e, x \in A$ satisfy*

$$0 \leq e \leq 1, \quad 0 \leq x \leq 1, \quad \text{and} \quad \|ex - x\| < \delta,$$

then $\|e^{1/2}x - x\| < \varepsilon$.

Proof. Choose a polynomial $p_0(\lambda) = \sum_{k=0}^n \beta_k \lambda^k$ with coefficients $\beta_k \in \mathbb{R}$ such that $|p_0(\lambda) - \lambda^{1/2}| < \frac{\varepsilon}{8}$ for all $\lambda \in [0, 1]$. Define

$$\alpha_0 = 0, \quad \alpha_1 = 1 - \sum_{k=2}^n \beta_k, \quad \alpha_2 = \beta_2, \quad \alpha_3 = \beta_3, \quad \dots, \quad \alpha_n = \beta_n.$$

Since $|p_0(0)| < \frac{\varepsilon}{8}$, we have $|\alpha_0 - \beta_0| < \frac{\varepsilon}{8}$. Since $|p_0(1) - 1| < \frac{\varepsilon}{8}$, we have

$$|\alpha_1 - \beta_1| = \left| 1 - \beta_1 - \sum_{k=2}^n \beta_k \right| \leq \left| 1 - \sum_{k=0}^n \beta_k \right| + |\beta_0| = |1 - p_0(1)| + |\beta_0| < \frac{2\varepsilon}{8}.$$

Therefore, for $\lambda \in [0, 1]$, the polynomial $p(\lambda) = \sum_{k=0}^n \alpha_k \lambda^k$ satisfies

$$|p(\lambda) - \lambda^{1/2}| \leq |\alpha_0 - \beta_0| + |\alpha_1 - \beta_1| \lambda + |p_0(\lambda) - \lambda^{1/2}| < \frac{\varepsilon}{8} + \frac{2\varepsilon}{8} + \frac{\varepsilon}{8} = \frac{\varepsilon}{2}.$$

Moreover $\alpha_0 = 0$ and $\sum_{k=0}^n \alpha_k = 1$.

Define

$$\delta = \frac{\varepsilon}{1 + 2 \sum_{k=1}^n k |\alpha_k|}.$$

Suppose now $\|ex - x\| < \delta$. An induction argument, using the estimate

$$\|e^k x - x\| \leq \|e^{k-1}\| \cdot \|ex - x\| + \|e^{k-1}x - x\|,$$

shows that for all $k \in \mathbb{Z}_{>0}$ we have $\|e^k x - x\| < k\delta$. Now, using $\sum_{k=1}^n \alpha_k x = x$ at the second step, we get

$$\begin{aligned} \|e^{1/2}x - x\| &\leq \|e^{1/2} - p(e)\| \cdot \|x\| + \|p(e)x - x\| \\ &\leq \|e^{1/2} - p(e)\| + \sum_{k=1}^n |\alpha_k| \cdot \|e^k x - x\| < \frac{\varepsilon}{2} + \sum_{k=0}^n k |\alpha_k| \delta < \varepsilon. \end{aligned}$$

This completes the proof. $\blacksquare$

Lemma 4.5. *Let A be a simple unital C^* -algebra, let $B \subset A$ be a large subalgebra of crossed product type (Definition 4.1), and let $C \subset B$ be the subalgebra of Definition 4.1. Assume $B \neq A$. Then for every $y \in B_+ \setminus \{0\}$ there is $z \in C_+ \setminus \{0\}$ such that $z \preceq_B y$.*

Proof. Choose any $a \in A$ such that $\text{dist}(a, B) > 2$. Definition 4.1 gives $c \in A$ and $g \in C$ such that

$$0 \leq g \leq 1, \quad \|c - a\| < 1, \quad (1 - g)c \in B, \quad \text{and} \quad g \preceq_B y.$$

The second condition implies that $c \notin B$, so the third condition implies that $g \neq 0$. Therefore the lemma is proved by taking $z = g$. $\blacksquare$

Theorem 4.6. *Let A be a stably finite simple unital C^* -algebra, and let $B \subset A$ be a large subalgebra of crossed product type (Definition 4.1). Then B is centrally large in A (Definition 3.1).*

We hope that finiteness of A is not necessary, but we do not know how to prove this. The difficulty is with condition (5) in the proof; when A is finite, this condition is, in effect, automatic.

We will use a weak (and obvious) variant of Proposition 3.8: we only restrict to a generating set to verify the approximate commutation condition.

Proof of Theorem 4.6. For a finite set T and $b_t \in A$ for $t \in T$, we take $\bigoplus_{t \in T} b_t \in M_{\text{card}(T)}(A)$ to be the diagonal matrix with diagonal entries b_t for $t \in T$. (The order won't matter in this proof.) In particular, $\bigoplus_{t \in T} b$ is a $\text{card}(T) \times \text{card}(T)$ diagonal matrix with all entries equal to b .

If $B = A$, then the conclusion is trivial. (We can always take $g = 0$.) We may therefore assume that $B \neq A$.

We must prove that for every $m \in \mathbb{Z}_{>0}$, $a_1, a_2, \dots, a_m \in A$, $\varepsilon > 0$, $x \in A_+$ with $\|x\| = 1$, and $y \in B_+ \setminus \{0\}$, there are $c_1, c_2, \dots, c_m \in A$ and $g \in B$ such that:

- (1) $0 \leq g \leq 1$.
- (2) For $j = 1, 2, \dots, m$ we have $\|c_j - a_j\| < \varepsilon$.
- (3) For $j = 1, 2, \dots, m$ we have $(1 - g)c_j \in B$.
- (4) $g \preceq_B y$ and $g \preceq_A x$.
- (5) $\|(1 - g)x(1 - g)\| > 1 - \varepsilon$.
- (6) For $j = 1, 2, \dots, m$ we have $\|ga_j - a_jg\| < \varepsilon$.

It is clearly equivalent to choose some fixed set $S \subset A$ which generates A as a C^* -algebra, and then to prove that for every $m \in \mathbb{Z}_{>0}$, $a_1, a_2, \dots, a_m \in A$, finite subset $F \subset S$, $\varepsilon > 0$, $x \in A_+$ with $\|x\| = 1$, and $y \in B_+ \setminus \{0\}$, there are $c_1, c_2, \dots, c_m \in A$ and $g \in B$ such that (1), (2), (3), (4), and (5) above hold, and such that in place of (6) the following holds:

- (7) For every $x \in F$ we have $\|gx - xg\| < \varepsilon$.

Now let $C \subset B$ and $G \subset A$ be as in Definition 4.1. We take the set S above to be the union of G and the unitary group of C . Then S is a set of unitaries which generates A as a C^* -algebra, and $uC u^* \subset C$ and $u^* C u \subset C$ for all $u \in S$.

Let $m \in \mathbb{Z}_{>0}$, let $a_1, a_2, \dots, a_m \in A$, let $F \subset S$ be finite, let $\varepsilon > 0$, let $x \in A_+$ satisfy $\|x\| = 1$, and let $y \in B_+ \setminus \{0\}$. Choose $n \in \mathbb{Z}_{>0}$ such that

$$(4.1) \quad \frac{1}{n} < \frac{\varepsilon}{8}.$$

Set

$$F_0 = \{1\}, \quad F_1 = F \cup F^* \cup \{1\}, \quad F_2 = F_1 F_1, \quad F_3 = F_1 F_2, \quad \dots, \quad F_n = F_1 F_{n-1}.$$

Set $N = \text{card}(F_1)$. Use Lemma 2.9 of [16] to choose $y_0 \in (\overline{x A x})_+ \setminus \{0\}$ such that:

- (8) If $g \in A_+$ satisfies $0 \leq g \leq 1$ and $g \preceq_A y_0$, then $\|(1 - g)x(1 - g)\| > 1 - \varepsilon$.

In particular,

$$(4.2) \quad y_0 \preceq_A x.$$

Proposition 4.2 implies that B is a large subalgebra of A , and Proposition 5.2 of [16] now implies that B is simple. Use Lemma 5.3 of [16] with $r = 1$, $a = y_0$, and $\varepsilon = 1$ to choose $y_1 \in B_+ \setminus \{0\}$ such that

$$(4.3) \quad y_1 \preceq_A y_0.$$

(We don't need part (3) of the conclusion of this lemma, and part (1) is only used to ensure that $y_1 \neq 0$.) Use simplicity of B and Lemma 2.6 of [16] to choose $y_2 \in B_+ \setminus \{0\}$ such that

$$(4.4) \quad y_2 \preceq_B y_1 \quad \text{and} \quad y_2 \preceq_B y.$$

Since B is simple, Lemma 2.4 of [16] provides a nonzero positive element $y_3 \in \overline{y_2 B y_2}$ such that the following condition holds:

- (9) There are $N^{n-1} + 1$ orthogonal positive elements in $\overline{y_2 B y_2}$, each Cuntz equivalent in B to y_3 .

Now use Lemma 4.5 to choose $z_0 \in C_+ \setminus \{0\}$ such that

$$(4.5) \quad z_0 \preceq_B y_3.$$

Then $u^* z_0 u \in C \subset B$ for all $u \in F_{n-1}$. Use Lemma 2.6 of [16] (applied to B) to choose $z_1 \in B_+ \setminus \{0\}$ such that

$$(4.6) \quad z_1 \preceq_B u^* z_0 u$$

for all $u \in F_{n-1}$. Then use Lemma 4.5 to choose $z \in C_+ \setminus \{0\}$ such that

$$(4.7) \quad z \preceq_B z_1.$$

Since A is simple, Lemma 2.4 of [16] provides a nonzero positive element $x_0 \in \overline{x A x}$ such that the following condition holds:

- (10) There are $\text{card}(F_{n-1})$ orthogonal positive elements in $\overline{x A x}$, each Cuntz equivalent in A to x_0 .

Apply Definition 4.1 with $a_1, a_2, \dots, a_m$, and ε as given, and with z in place of y and x_0 in place of x . Let $c_1, c_2, \dots, c_m \in A$ and $g_0 \in C$ be the resulting elements. In particular,

$$(4.8) \quad g_0 \preceq_B z \quad \text{and} \quad g_0 \preceq_A x_0.$$

Choose $\varepsilon_0 > 0$ so small that the following three things happen:

- (11) $\varepsilon_0 < \frac{\varepsilon}{8}$.
(12) The choice $2\varepsilon_0$ can be used for δ in Lemma 2.5 when ε there is taken to be $\frac{\varepsilon}{8}$ and when $f(x) = x^{1/2}$.
(13) The choice ε_0 can be used for δ in Lemma 4.4 when ε there is taken to be $\frac{\varepsilon}{8}$.

For $k = 0, 1, \dots, n$ and all $u \in F_k$, we have $u g_0 u^* \in C$. For $k = 0, 1, \dots, n$, let $D_k \subset B$ be the hereditary subalgebra of B generated by all $u g_0 u^*$ for $u \in F_k$. Then $u D_k u^* \subset D_{k+1}$ for $u \in F_1$ and $k = 0, 1, \dots, n-1$. By induction, choose

$$g_1 \in D_1, \quad g_2 \in D_2, \quad \dots, \quad g_n \in D_n$$

such that

$$(4.9) \quad 0 \leq g_k \leq 1 \quad \text{and} \quad \|g_k u g_l u^* - u g_l u^*\| < \varepsilon_0$$

for

$$k = 1, 2, \dots, n, \quad u \in F_1, \quad \text{and} \quad l = 0, 1, \dots, k-1.$$

Now define

$$(4.10) \quad r_0 = \left(1 - \frac{1}{n} \sum_{k=1}^{n-1} g_k\right)^{1/2}, \quad r = r_0^* r_0, \quad \text{and} \quad g = 1 - r.$$

We verify conditions (1), (2), (3), (4), (5), and (7).

For (1), we clearly have $\|r_0\| \leq 1$, so $0 \leq r \leq 1$, whence $0 \leq g \leq 1$.

Condition (2) follows from the choice of $c_1, c_2, \dots, c_m$.

We verify condition (3). Corollary 2.7 implies that $(1 - g_0)^{1/2} c_j \in B$ for $j = 1, 2, \dots, m$. Since $(1 - g_0)^{1/2} \in B$ and $1 - \frac{1}{n} \sum_{k=1}^{n-1} g_k \in B$, it now follows that $r c_j \in B$ for $j = 1, 2, \dots, m$. This is (3).

Now we prove (4) and (5). Set

$$(4.11) \quad s = 1 - (1 - g_0)^{1/2} \quad \text{and} \quad h = \frac{1}{n} \sum_{k=0}^{n-1} g_k.$$

Then $s = f(g_0)$ for the continuous function $f: [0, 1] \rightarrow [0, 1]$ given by $f(\lambda) = 1 - (1 - \lambda)^{1/2}$. Since $f(0) = 0$ and $g_0 \in D_0$, it follows that $s \in D_0 \subset D_{n-1}$. Also $h - \frac{1}{n}g_0 \in D_{n-1}$. Therefore

$$\begin{aligned} g &= 1 - r = 1 - (1 - s)[1 - (h - \frac{1}{n}g_0)](1 - s) \\ &= 2s + h - \frac{1}{n}g_0 - s(h - \frac{1}{n}g_0) - s^2 - (h - \frac{1}{n}g_0)s + s(h - \frac{1}{n}g_0)s \in D_{n-1}. \end{aligned}$$

Now $d = \sum_{u \in F_{n-1}} u g_0 u^*$ satisfies $\overline{d D_{n-1} d} = D_{n-1}$. In B , using Proposition 2.7(i) of [12] at the first step and (4.8) at the second step, we now get

$$g \preceq_B \sum_{u \in F_{n-1}} u g_0 u^* \preceq_A \bigoplus_{u \in F_{n-1}} u z u^*.$$

For $u \in F_{n-1}$, we have $z \preceq_B u^* z_0 u$ by (4.7) and (4.6), so $u z u^* \preceq_A z_0$. Thus, using (4.5) at the last step,

$$g \preceq_A \bigoplus_{u \in F_{n-1}} u z u^* \preceq_A \bigoplus_{u \in F_{n-1}} z_0 \preceq_B \bigoplus_{u \in F_{n-1}} y_3.$$

Now

$$y_3 \oplus \bigoplus_{u \in F_{n-1}} y_3 \preceq_B y_2$$

by (9). Thus $y_3 \oplus g \preceq_A y_2$. Corollary 5.8 of [16] implies that B is stably large in A . So we can apply Lemma 6.5 of [16], with $a = g$, $b = y_2$, $c = \bigoplus_{u \in F_{n-1}} y_3$, and $x = y_3$, to get $g \preceq_B y_2$. Thus (4.4) gives $g \preceq_B y_2 \preceq_B y$. By (4.4) and (4.3) we also get $g \preceq_B y_2 \preceq_B y_1 \preceq_A y_0$, so (8) implies that $\|(1 - g)x(1 - g)\| > 1 - \varepsilon$, and (4.2) implies $g \preceq_A x$.

Finally, we prove (7). We first claim that

$$(4.12) \quad \left\| \left[1 - g_0, 1 - \frac{1}{n} \sum_{k=1}^{n-1} g_k \right] \right\| < 2\varepsilon_0.$$

It is enough to prove that

$$\left\| \left[g_0, \frac{1}{n} \sum_{k=1}^{n-1} g_k \right] \right\| < 2\varepsilon_0.$$

The claim thus follows by using (4.9) to get, for $k = 1, 2, \dots, n-1$,

$$\|g_k g_0 - g_0 g_k\| \leq \|g_k g_0 - g_0\| + \|g_0 - g_0 g_k\| < \varepsilon_0 + \varepsilon_0 = 2\varepsilon_0.$$

Using (12) and (4.12), we get

$$\left\| \left[(1 - g_0)^{1/2}, 1 - \frac{1}{n} \sum_{k=1}^{n-1} g_k \right] \right\| < \frac{\varepsilon}{8}.$$

Therefore

$$(4.13) \quad \left\| r - (1 - g_0) \left(1 - \frac{1}{n} \sum_{k=1}^{n-1} g_k \right) \right\| < \frac{\varepsilon}{8}.$$

Recall the definition of h in (4.11). We now claim that

$$(4.14) \quad \|h - g\| < \frac{\varepsilon}{4}.$$

To prove this, we use (4.13) and $g = 1 - r$ (from (4.10)) at the first step, $1 \in F_1$ and (4.9) at the third last step, and (11) at the second last step, to estimate

$$\begin{aligned} \|h - g\| &< \frac{\varepsilon}{8} + \left\| 1 - (1 - g_0) \left(1 - \frac{1}{n} \sum_{k=1}^{n-1} g_k \right) - h \right\| \\ &= \frac{\varepsilon}{8} + \left\| g_0 + \frac{1}{n} \sum_{k=1}^{n-1} g_k - \frac{1}{n} \sum_{k=1}^{n-1} g_0 g_k - \frac{1}{n} \sum_{k=0}^{n-1} g_k \right\| \\ &= \frac{\varepsilon}{8} + \left\| \left(\frac{n-1}{n} \right) g_0 - \frac{1}{n} \sum_{k=1}^{n-1} g_0 g_k \right\| \leq \frac{\varepsilon}{8} + \frac{1}{n} \sum_{k=1}^{n-1} \|g_0 - g_0 g_k\| \\ &< \frac{\varepsilon}{8} + \varepsilon_0 \leq \frac{\varepsilon}{8} + \frac{\varepsilon}{8} = \frac{\varepsilon}{4}. \end{aligned}$$

The claim is proved.

We next claim that $\|uhu^* - h\| \leq \frac{\varepsilon}{2}$ for $u \in F$. Set $g_{-1} = 0$ and define

$$e = \frac{1}{n} \sum_{k=0}^{n-1} u g_k^{1/2} u^* g_{k-1} u g_k^{1/2} u^* \quad \text{and} \quad f = \frac{1}{n} \sum_{k=0}^{n-1} g_{k+1}^{1/2} u g_k u^* g_{k+1}^{1/2}.$$

We will apply Lemma 4.3 with

$$a = \frac{1}{n} \sum_{k=0}^{n-2} g_k, \quad b = \frac{1}{n} \sum_{k=0}^n g_k, \quad d = h, \quad r = uhu^*, \quad x = e, \quad y = f,$$

and with $\frac{2}{n}$ in place of ε and $\frac{\varepsilon}{4}$ in place of ρ . We verify its hypotheses. The relation

$$\frac{1}{n} \sum_{k=0}^{n-2} g_k \leq h \leq \frac{1}{n} \sum_{k=0}^n g_k$$

is clear. We also have

$$\left\| \frac{1}{n} \sum_{k=0}^n g_k - \frac{1}{n} \sum_{k=0}^{n-2} g_k \right\| \leq \frac{1}{n} (\|g_{n-1}\| + \|g_n\|) \leq \frac{2}{n}.$$

The relation $f \leq \frac{1}{n} \sum_{k=0}^n g_k$ follows from $g_0 \geq 0$ and

$$g_{k+1}^{1/2} u g_k u^* g_{k+1}^{1/2} \leq \|u g_k u^*\| g_{k+1} \leq g_{k+1}.$$

The proof of the inequality $e \leq uhu^*$ is similar.

We now check that

$$(4.15) \quad \left\| e - \frac{1}{n} \sum_{k=0}^{n-2} g_k \right\| \leq \frac{\varepsilon}{4}.$$

For $k = 1, 2, \dots, n-1$, use (4.9) and (13) at the second step to get

$$\|u g_k^{1/2} u^* g_{k-1} - g_{k-1}\| = \|g_k^{1/2} u^* g_{k-1} u - u^* g_{k-1} u\| < \frac{\varepsilon}{8}.$$

Taking adjoints gives also

$$\|g_{k-1} u g_k^{1/2} u^* - g_{k-1}\| < \frac{\varepsilon}{8}.$$

So

$$\|ug_k^{1/2}u^*g_{k-1}ug_k^{1/2}u^* - g_{k-1}\| < \frac{\varepsilon}{4}.$$

Since $g_{-1} = 0$, we therefore have

$$\begin{aligned} \left\| e - \frac{1}{n} \sum_{k=0}^{n-2} g_k \right\| &= \left\| \frac{1}{n} \sum_{k=1}^{n-1} ug_k^{1/2}u^*g_{k-1}ug_k^{1/2}u^* - \frac{1}{n} \sum_{k=1}^{n-1} g_{k-1} \right\| \\ &\leq \frac{1}{n} \sum_{k=1}^{n-1} \|ug_k^{1/2}u^*g_{k-1}ug_k^{1/2}u^* - g_{k-1}\| < \frac{\varepsilon}{4}. \end{aligned}$$

This is (4.15).

Similar reasoning to that of the previous paragraph gives

$$\|g_{k+1}^{1/2}ug_ku^*g_{k+1}^{1/2} - ug_ku^*\| < \frac{\varepsilon}{4}$$

for $k = 1, 2, \dots, n$, so that

$$\|f - uhu^*\| \leq \frac{1}{n} \sum_{k=0}^{n-1} \|g_{k+1}^{1/2}ug_ku^*g_{k+1}^{1/2} - ug_ku^*\| < \frac{\varepsilon}{4}.$$

This completes the verification of the hypotheses of Lemma 4.3. Applying this lemma, and using (4.1) at the second step, we get

$$\|uhu^* - h\| \leq \frac{2}{n} + \frac{\varepsilon}{4} < \frac{\varepsilon}{2}.$$

Combining this estimate with (4.14), for $u \in F$ we now have

$$\|[g, u]\| = \|g - ugu^*\| \leq \|h - uhu^*\| + 2\|g - h\| < \frac{\varepsilon}{2} + 2\left(\frac{\varepsilon}{4}\right) = \varepsilon.$$

This completes the proof of condition (7), and the proof of the theorem. $\blacksquare$


5. REPLACING THE MATRIX DECOMPOSITION

One frequently used technique in C*-algebras with many projections is decomposing the identity into orthogonal projections in order to create a matrix decomposition of some element of interest. Lemma 5.4 below creates a decomposition of the identity into three positive elements z_1 , z_2 , and z_3 (which are not mutually orthogonal). These elements are used to simulate a 3×3 matrix decomposition of an element of the C*-algebra. It is helpful to think of the following picture.



Lemma 5.2 is a technical lemma using two related decompositions of the identity into three positive elements and simulating a 5×5 matrix decomposition

of an element of the C^* -algebra. Lemma 5.2 is used to prove Lemma 5.4. The following is the motivational picture.



The next lemma is used repeatedly, but often implicitly, in the proof of Lemma 5.2.

Lemma 5.1. *If y and z are orthogonal positive elements of a C^* -algebra A and $w \in \overline{Ay}$ and $x \in \overline{zA}$, then $wx = 0$.*

Proof. The statement is true if $w \in Ay$ and $x \in zA$. Take limits. ■

Lemma 5.2. *Let A be a finite unital C^* -algebra and let $a \in A$. Let $B \subset A$ be a unital subalgebra with $\text{tsr}(B) = 1$. Suppose there exist positive elements $b_1, b_2, b_3, c_1, c_2, c_3 \in B$ such that:*

- (1) $b_1 + b_2 + b_3 = 1$.
- (2) $c_1 + c_2 + c_3 = 1$.
- (3) $C^*(b_1, b_2, b_3, c_1, c_2, c_3)$ is commutative.
- (4) $b_1 c_1 = c_1$.
- (5) $b_3 c_3 = c_3$.
- (6) $b_2 c_2 = b_2$.
- (7) $c_1 b_2 = c_3 b_2 = 0$.
- (8) $b_1 b_3 = 0$.
- (9) $c_2 a c_2 \in \overline{c_2 B c_2}$.
- (10) $b_1 a = a b_3 = 0$.

Then for all $\varepsilon > 0$ there exists an invertible element $y \in A$ such that $\|a - y\| < \varepsilon$.

Proof. The relations in the hypotheses imply:

$$(5.1) \quad c_1 b_3 = c_1 b_1 b_3 = 0.$$

$$(5.2) \quad c_1 c_3 = c_1 b_3 c_3 = 0.$$

$$(5.3) \quad b_1 c_3 = b_1 b_3 c_3 = 0.$$

From (1) we get

$$(5.4) \quad 1 = c_1 + (b_1 - c_1) + b_2 + (b_3 - c_3) + c_3.$$

We wish to use this decomposition of the identity to decompose a . Therefore, make the following definitions:

$$(5.5) \quad a_{3,1} = b_2 a c_1.$$

$$(5.6) \quad a_{3,2} = b_2 a (b_1 - c_1).$$

$$(5.7) \quad a_{3,3} = b_2 a b_2.$$

$$(5.8) \quad a_{4,1} = (b_3 - c_3)ac_1.$$

$$(5.9) \quad a_{4,2} = (b_3 - c_3)a(b_1 - c_1).$$

$$(5.10) \quad a_{4,3} = (b_3 - c_3)ab_2.$$

$$(5.11) \quad a_{5,1} = c_3ac_1.$$

$$(5.12) \quad a_{5,2} = c_3a(b_1 - c_1).$$

$$(5.13) \quad a_{5,3} = c_3ab_2.$$

Then

$$(5.14) \quad \sum_{j=3}^5 \sum_{k=1}^3 a_{j,k} = a$$

by (5.4) and the hypotheses (10), (4), and (5).

We have, using (4) at the first step, (2) at the second step, and (5.3) at the third step,

$$b_1 - c_1 = b_1(1 - c_1) = b_1(c_2 + c_3) = b_1c_2.$$

Similarly $b_3 - c_3 = b_3c_2$. Also using (6), it follows that

$$a_{3,2}, a_{3,3}, a_{4,2}, a_{4,3} \in \overline{c_2Bc_2};$$

in fact, $a_{4,3} \in \overline{(b_3 - c_3)Bb_2}$. Since $b_2c_2 = b_2$, we have $b_2ab_2 \in \overline{b_2Bb_2} \subset \overline{c_2Bc_2}$. Furthermore, since B has stable rank one, $\overline{b_2Bb_2}$ and $\overline{c_2Bc_2}$ also have stable rank one.

Set

$$\delta = \min \left(\frac{\varepsilon}{6\|a_{3,1} + a_{4,1} + a_{5,1}\|}, \sqrt{\frac{\varepsilon}{6}}, \frac{\varepsilon}{3} \right).$$

Since $a_{3,3} \in \overline{b_2Bb_2}$, there is an invertible element $t_0 \in \overline{b_2Bb_2} + \mathbb{C} \cdot 1_A$ with

$$(5.15) \quad \|t_0 - a_{3,3}\| < \delta.$$

Then there are $t_1, t_2 \in \overline{b_2Bb_2}$ and $\lambda_1 \in \mathbb{C}$ such that $t_0 = t_1 + \lambda_1 1_A$ and $t_0^{-1} = t_2 + \lambda_1^{-1} 1_A$.

We claim that

$$(5.16) \quad ((a_{3,1} + a_{4,1} + a_{5,1})t_0^{-1})^2 = 0.$$

To prove the claim, we first observe that, using the hypothesis (7) and Lemma 5.1 at the second step, we get

$$(a_{3,1} + a_{4,1} + a_{5,1})t_0 = (a_{3,1} + a_{4,1} + a_{5,1})(t_1 + \lambda_1) = (a_{3,1} + a_{4,1} + a_{5,1})\lambda_1.$$

Therefore

$$a_{3,1} + a_{4,1} + a_{5,1} = (a_{3,1} + a_{4,1} + a_{5,1})t_0t_0^{-1} = (a_{3,1} + a_{4,1} + a_{5,1})\lambda_1t_0^{-1}.$$

Using (5.1), (5.2), and the hypothesis (7) at the second step, this implies

$$((a_{3,1} + a_{4,1} + a_{5,1})t_0^{-1})^2 = (a_{3,1} + a_{4,1} + a_{5,1})^2\lambda_1^{-2} = 0.$$

The claim is proved.

Now we compute

$$\begin{aligned}
 (5.17) \quad & (a_{3,1} + a_{4,1} + a_{5,1} + t_0) t_0^{-1} [1 - (a_{3,1} + a_{4,1} + a_{5,1}) t_0^{-1}] \\
 &= (a_{3,1} + a_{4,1} + a_{5,1}) t_0^{-1} - ((a_{3,1} + a_{4,1} + a_{5,1}) t_0^{-1})^2 \\
 &\quad + 1 - (a_{3,1} + a_{4,1} + a_{5,1}) t_0^{-1} \\
 &= 1.
 \end{aligned}$$

Because A is finite, every one sided invertible element is invertible. So $a_{3,1} + a_{4,1} + a_{5,1} + t_0$ is invertible, with

$$(5.18) \quad (a_{3,1} + a_{4,1} + a_{5,1} + t_0)^{-1} = t_0^{-1} [1 - (a_{3,1} + a_{4,1} + a_{5,1}) t_0^{-1}].$$

Using our expression for t_0^{-1} in the first step and using Lemma 5.1 twice in the second step, once with $y = c_1$ and $z = b_2$ and once with $y = c_1$ and $z = b_3$ (using (5.1)), we compute

$$(5.19) \quad (a_{3,1} + a_{4,1} + a_{5,1}) t_0^{-1} b_3 = (a_{3,1} + a_{4,1} + a_{5,1}) (t_2 + \lambda_1^{-1}) b_3 = 0.$$

Similarly,

$$(5.20) \quad (a_{3,1} + a_{4,1} + a_{5,1}) t_0^{-1} b_2 = 0.$$

From (5.19) and the hypothesis (5), we get

$$(5.21) \quad (a_{3,1} + a_{4,1} + a_{5,1}) t_0^{-1} c_3 = 0 \quad \text{and} \quad (a_{3,1} + a_{4,1} + a_{5,1}) t_0^{-1} (b_3 - c_3) = 0.$$

Next we multiply, using (5.18) at the first step, and at the last step using $t_2 \in \overline{b_2 B b_2}$ together with the hypothesis (7) on the terms $t_2 a_{5,2}$ and $t_2 a_{5,3}$, and using (5.20) and (5.21) on the terms starting with t_0^{-1} :

$$\begin{aligned}
 (5.22) \quad & t_0^{-1} [1 - (a_{3,1} + a_{4,1} + a_{5,1}) t_0^{-1}] \\
 &\quad \cdot (a_{3,1} + a_{3,2} + t_0 + a_{4,1} + a_{4,2} + a_{4,3} + a_{5,1} + a_{5,2} + a_{5,3}) \\
 &= 1 + t_0^{-1} [1 - (a_{3,1} + a_{4,1} + a_{5,1}) t_0^{-1}] \\
 &\quad \cdot (a_{3,2} + a_{4,2} + a_{4,3} + a_{5,2} + a_{5,3}) \\
 &= 1 + [t_2 + \lambda_1^{-1} - t_0^{-1} (a_{3,1} + a_{4,1} + a_{5,1}) t_0^{-1}] \\
 &\quad \cdot (a_{3,2} + a_{4,2} + a_{4,3} + a_{5,2} + a_{5,3}) \\
 &= 1 + t_2 a_{3,2} + t_2 a_{4,2} + t_2 a_{4,3} + t_2 a_{5,2} + t_2 a_{5,3} \\
 &\quad + \lambda_1^{-1} a_{3,2} + \lambda_1^{-1} a_{4,2} + \lambda_1^{-1} a_{4,3} + \lambda_1^{-1} a_{5,2} + \lambda_1^{-1} a_{5,3} \\
 &\quad - t_0^{-1} (a_{3,1} + a_{4,1} + a_{5,1}) t_0^{-1} a_{3,2} \\
 &\quad - t_0^{-1} (a_{3,1} + a_{4,1} + a_{5,1}) t_0^{-1} (a_{4,2} + a_{4,3}) \\
 &\quad - t_0^{-1} (a_{3,1} + a_{4,1} + a_{5,1}) t_0^{-1} (a_{5,2} + a_{5,3}) \\
 &= 1 + t_2 a_{3,2} + t_2 a_{4,2} + t_2 a_{4,3} + \lambda_1^{-1} a_{3,2} \\
 &\quad + \lambda_1^{-1} a_{4,2} + \lambda_1^{-1} a_{4,3} + \lambda_1^{-1} a_{5,2} + \lambda_1^{-1} a_{5,3}.
 \end{aligned}$$

Set $t_3 = t_2 a_{4,3} + 1$. Since $t_2 \in \overline{b_2 B b_2}$ and $a_{4,3} \in \overline{(b_3 - c_3) B b_2}$, we have $t_3 \in \overline{b_2 B b_2} + \mathbb{C} \cdot 1_A$. Thus there is an invertible element $t_4 \in \overline{b_2 B b_2} + \mathbb{C} \cdot 1_A$ with

$$(5.23) \quad \|t_4 - t_3\| < \delta.$$

So there are $t_5, t_6 \in \overline{b_2 B b_2}$ and $\lambda_5 \in \mathbb{C}$ such that $t_4 = t_5 + \lambda_5 1_A$ and $t_4^{-1} = t_6 + \lambda_5^{-1} 1_A$.

An argument similar to that for Equation (5.16), using (5.2), (5.3), and the hypothesis (7) gives $(t_4^{-1}(a_{5,2} + a_{5,3}))^2 = 0$. Define

$$w = \lambda_1^{-1}a_{5,2} + \lambda_1^{-1}a_{5,3}.$$

Thus

$$(5.24) \quad (t_4^{-1}w)^2 = 0.$$

A computation similar to the one for Equation (5.17) shows that $w + t_4$ is invertible, with

$$(5.25) \quad (w + t_4)^{-1} = (1 - t_4^{-1}w)t_4^{-1}.$$

Applying Lemma 5.1, and using (5.3), the hypothesis (7), and the relations

$$t_6 \in \overline{b_2 B b_2}, \quad b_1 \in \overline{b_1 B b_1}, \quad \text{and} \quad a_{5,2} + a_{5,3} \in \overline{c_3 A},$$

we get

$$(5.26) \quad b_1 t_4^{-1}w = b_1 t_6 w + b_1 \lambda_5^{-1}w = 0.$$

Similarly,

$$(5.27) \quad b_2 t_4^{-1}w = 0.$$

Hypothesis (4) and the relation (5.26) also imply that

$$(5.28) \quad c_1 t_4^{-1}w = 0 \quad \text{and} \quad (b_1 - c_1)t_4^{-1}w = 0.$$

Define

$$\begin{aligned} t_7 = 1 &+ t_2 a_{3,2} t_6 + t_2 a_{4,2} t_6 + \lambda_1^{-1} a_{3,2} t_6 + \lambda_1^{-1} a_{4,2} t_6 + \lambda_1^{-1} a_{4,3} t_6 \\ &+ \lambda_5^{-1} t_2 a_{3,2} + \lambda_5^{-1} t_2 a_{4,2} + \lambda_1^{-1} \lambda_5^{-1} a_{3,2} + \lambda_1^{-1} \lambda_5^{-1} a_{4,2} + \lambda_1^{-1} \lambda_5^{-1} a_{4,3}. \end{aligned}$$

One checks that $t_7 \in \overline{c_2 B c_2} + \mathbb{C} \cdot 1_A$, which has stable rank one by hypothesis. Thus there exists an invertible element $t_8 \in \overline{c_2 B c_2} + \mathbb{C} \cdot 1_A$ such that

$$(5.29) \quad \|t_8 - t_7\| < \frac{\delta}{1 + \|w\|}.$$

Set

$$y = (a_{3,1} + a_{4,1} + a_{5,1} + t_0) t_8 (w + t_4).$$

Then y is invertible by (5.18) and (5.25).

We claim that $\|a - y\| < \varepsilon$. To prove the claim, we begin with the following computation. In the first step we multiply out, cancel the terms $\pm w t_4^{-1}$, and use (5.24) to see that $w t_4^{-1} w t_4^{-1} = 0$. The second step uses (5.27) and (5.28) to see that the terms involving $t_4^{-1} w t_4^{-1}$ vanish. The third step is obtained by substituting $t_4^{-1} = t_6 + \lambda_5^{-1} 1_A$ and comparing with the definition of t_7 . We have:

$$\begin{aligned} (5.30) \quad & (t_2 a_{3,2} + t_2 a_{4,2} + t_4 + \lambda_1^{-1} a_{3,2} + \lambda_1^{-1} a_{4,2} + \lambda_1^{-1} a_{4,3} + w) \cdot [1 - t_4^{-1}w] t_4^{-1} \\ &= 1 + t_2 a_{3,2} t_4^{-1} + t_2 a_{4,2} t_4^{-1} + \lambda_1^{-1} a_{3,2} t_4^{-1} + \lambda_1^{-1} a_{4,2} t_4^{-1} + \lambda_1^{-1} a_{4,3} t_4^{-1} \\ &\quad - t_2 a_{3,2} t_4^{-1} w t_4^{-1} - t_2 a_{4,2} t_4^{-1} w t_4^{-1} \\ &\quad - \lambda_1^{-1} a_{3,2} t_4^{-1} w t_4^{-1} - \lambda_1^{-1} a_{4,2} t_4^{-1} w t_4^{-1} - \lambda_1^{-1} a_{4,3} t_4^{-1} w t_4^{-1} \\ &= 1 + t_2 a_{3,2} t_4^{-1} + t_2 a_{4,2} t_4^{-1} + \lambda_1^{-1} a_{3,2} t_4^{-1} + \lambda_1^{-1} a_{4,2} t_4^{-1} + \lambda_1^{-1} a_{4,3} t_4^{-1} \\ &= t_7. \end{aligned}$$

Next, using (5.25) at the first step, (5.30) at the second step, and the choices of t_4 and t_8 at the third step, we get

(5.31)

$$\begin{aligned}
& \left\| 1 + t_2 a_{3,2} + t_2 a_{4,2} + t_2 a_{4,3} + \lambda_1^{-1} a_{3,2} + \lambda_1^{-1} a_{4,2} + \lambda_1^{-1} a_{4,3} + w - t_8(w + t_4) \right\| \\
& \leq \left\| 1 + t_2 a_{4,3} - t_4 \right\| \\
& \quad + \left\| (t_2 a_{3,2} + t_2 a_{4,2} + t_4 + \lambda_1^{-1} a_{3,2} + \lambda_1^{-1} a_{4,2} + \lambda_1^{-1} a_{4,3} + w) \right. \\
& \quad \cdot (1 - t_4^{-1} w) t_4^{-1} (w + t_4) \\
& \quad \left. - t_8(w + t_4) \right\| \\
& \leq \left\| 1 + t_2 a_{4,3} - t_4 \right\| + \left\| t_7 - t_8 \right\| \cdot \left\| w + t_4 \right\| \\
& \leq \delta + \left(\frac{\delta}{\left\| w + t_4 \right\| + 1} \right) \left\| w + t_4 \right\| < 2\delta.
\end{aligned}$$

Then we compute as follows, using (5.14) and the definitions of y and w at the first step, (5.17) at the second step, (5.22) and the definition of w at the third step, and (5.31) at the fourth step,

$$\begin{aligned}
\|a - y\| &= \left\| a_{3,1} + a_{3,2} + a_{3,3} + a_{4,1} + a_{4,2} + a_{4,3} + a_{5,1} + \lambda_1 w \right. \\
&\quad \left. - (a_{3,1} + a_{4,1} + a_{5,1} + t_0) t_8(w + t_4) \right\| \\
&\leq \left\| a_{3,3} - t_0 \right\| \\
&\quad + \left\| (a_{3,1} + a_{4,1} + a_{5,1} + t_0) t_0^{-1} [1 - (a_{3,1} + a_{4,1} + a_{5,1}) t_0^{-1}] \right. \\
&\quad \cdot (a_{3,1} + a_{3,2} + t_0 + a_{4,1} + a_{4,2} + a_{4,3} + a_{5,1} + \lambda_1 w) \\
&\quad \left. - (a_{3,1} + a_{4,1} + a_{5,1} + t_0) t_8(w + t_4) \right\| \\
&\leq \delta + \left\| a_{3,1} + a_{4,1} + a_{5,1} + t_0 \right\| \\
&\quad \cdot \left\| 1 + t_2 a_{3,2} + t_2 a_{4,2} + t_2 a_{4,3} + \lambda_1^{-1} a_{3,2} + \lambda_1^{-1} a_{4,2} + \lambda_1^{-1} a_{4,3} \right. \\
&\quad \left. + w - t_8(w + t_4) \right\| \\
&\leq \delta + 2\delta \left\| a_{3,1} + a_{4,1} + a_{5,1} + t_0 \right\| \\
&\leq \delta + 2\delta \left\| a_{3,1} + a_{4,1} + a_{5,1} + a_{3,3} \right\| + 2\delta \left\| a_{3,3} - t_0 \right\| < \frac{\varepsilon}{3} + \frac{2\varepsilon}{6} + 2\delta^2 \leq \varepsilon.
\end{aligned}$$

This proves the claim and completes the proof. ■

Lemma 5.3. *Let A be a C^* -algebra and let $B \subset A$ be a subalgebra. Suppose $a \in A$ and $b \in B_+$ satisfy $bab \in B$. Let $f: \text{sp}(b) \rightarrow [0, \infty)$ be a continuous function vanishing at zero. Then $f(b)af(b) \in \overline{f(b)Bf(b)}$.*

Proof. We have $b^{1/n}ab^{1/n} \in B$ by a standard polynomial approximation argument. Also, for all $k \in \mathbb{Z}_{>0}$ we have $\lim_{n \rightarrow \infty} b^k b^{1/n} = b^k$. Therefore polynomial approximation gives $\lim_{n \rightarrow \infty} f(b)b^{1/n} = f(b)$. Similarly $\lim_{n \rightarrow \infty} b^{1/n}f(b) = f(b)$. The result follows. ■

Lemma 5.4. *Let A be a finite unital C^* -algebra and let $a \in A$. Let $B \subset A$ be a unital subalgebra with $\text{tsr}(B) = 1$. Suppose there exist $z_1, z_2, z_3 \in B$ such that:*

- (1) $z_1, z_2, z_3 \geq 0$.
- (2) $z_1 + z_2 + z_3 = 1$.
- (3) $z_1 z_3 = 0$.

- (4) $z_1 a = 0$.
- (5) $a z_3 = 0$
- (6) $z_2 a z_2 \in B$.

Then for all $\varepsilon > 0$ there exists an invertible element $y \in A$ such that $\|y - a\| < \varepsilon$.

Proof. We verify the hypotheses of Lemma 5.2. Let $f, h: [0, 1] \rightarrow [0, 1]$ be defined by the formulas

$$f(\lambda) = \begin{cases} 0 & \lambda \in [0, \frac{1}{2}] \\ 2\lambda - 1 & \lambda \in [\frac{1}{2}, 1] \end{cases} \quad \text{and} \quad h(\lambda) = \begin{cases} 2\lambda & \lambda \in [0, \frac{1}{2}] \\ 1 & \lambda \in [\frac{1}{2}, 1] \end{cases}.$$

Then $fh = f$.

Define positive elements $b_1, b_2, b_3, c_1, c_2, c_3 \in A$ by

$$b_1 = h(z_1), \quad b_2 = f(z_2), \quad b_3 = h(z_3), \quad c_1 = f(z_1), \quad c_2 = h(z_2), \quad c_3 = f(z_3).$$

Set $D = C^*(z_1, z_2, z_3)$. Since $D = C^*(1, z_1, z_3)$ and $z_1 z_3 = 0$, we see that D is commutative. So there is a compact metric space X and an isomorphism $D \cong C(X)$, which we treat as an identification. It also follows that $C^*(b_1, b_2, b_3, c_1, c_2, c_3) \subset D$ is commutative. We have $c_2 a c_2 \in \overline{c_2 B c_2}$ by Lemma 5.3.

Identifying $D = C(X)$ as above, for $j = 1, 2, 3$ set

$$U_j = \{x \in X : z_j(x) > 0\} \quad \text{and} \quad V_j = \{x \in X : z_j(x) > \frac{1}{2}\}.$$

From the definitions of f and h , we see that if $\lambda_1, \lambda_2 \in [0, 1]$ satisfy $\lambda_1 + \lambda_2 = 1$, then $f(\lambda_1) + h(\lambda_2) = 1$. Since $U_1 \cap U_3 = \emptyset$ and $z_1 + z_2 + z_3 = 1$, it is now easily checked that

$$b_1 + b_2 + b_3 = 1, \quad c_1 + c_2 + c_3 = 1, \quad \text{and} \quad b_1 b_3 = 0.$$

The relation $fh = f$ implies

$$b_1 c_1 = c_1, \quad b_2 c_2 = b_2, \quad \text{and} \quad b_3 c_3 = c_3.$$

For $x \in X$, we have $c_1(x) \neq 0$ if and only if $x \in V_1$ and $b_2(x) \neq 0$ if and only if $x \in V_2$. Clearly $V_1 \cap V_2 = \emptyset$, so $c_1 b_2 = 0$. Similarly $c_3 b_2 = 0$. We have $b_1 a = 0$ by Lemma 5.1 and similarly $a b_3 = 0$. This completes the verification of the hypotheses of Lemma 5.2. ■

6. CENTRALLY LARGE SUBALGEBRAS AND STABLE RANK ONE

In this section we will prove the main theorem (Theorem 6.3), which says that a C^* -algebra with a centrally large subalgebra which has stable rank one must itself have stable rank one. If the subalgebra also has real rank zero, we further show that the containing algebra does as well.

The proof of Theorem 6.3 proceeds via two technical lemmas. The first is a version of the definition of a centrally large subalgebra in which the element g is replaced by a tower of elements.

Lemma 6.1. *Let A be an infinite dimensional simple separable unital C^* -algebra, and let $B \subset A$ be a centrally large subalgebra. Then for all $m, N \in \mathbb{Z}_{>0}$, $a_1, a_2, \dots, a_m \in A$, $\varepsilon > 0$, $x \in A_+$ with $\|x\| = 1$, and $y \in B_+ \setminus \{0\}$, there are $c_1, c_2, \dots, c_m \in A$ and $g_0, g_1, \dots, g_N \in B$ such that:*

- (1) $0 \leq g_n \leq 1$ for $n = 0, 1, \dots, N$ and $g_{n-1} g_n = g_n$ for $n = 1, 2, \dots, N$.
- (2) For $j = 1, 2, \dots, m$ we have $\|c_j - a_j\| < \varepsilon$.
- (3) For $j = 1, 2, \dots, m$ and $n = 0, 1, \dots, N$, we have $(1 - g_n) c_j \in B$.

- (4) For $n = 0, 1, \dots, N$, we have $g_n \preceq_B y$ and $g_n \preceq_A x$.
- (5) For $n = 0, 1, \dots, N$, we have $\|(1 - g_n)x(1 - g_n)\| > 1 - \varepsilon$.
- (6) For $j = 1, 2, \dots, m$ and $n = 0, 1, \dots, N$, we have $\|g_n a_j - a_j g_n\| < \varepsilon$.

As we will see in the proof, it is enough to take $n = 0$ in (4) and (5) and $n = N$ in (3).

Proof of Lemma 6.1. Without loss of generality $\|a_j\| \leq 1$ for $j = 1, 2, \dots, m$ and $\varepsilon \leq 1$.

Define continuous functions $k_0, k_1, \dots, k_N: [0, 1] \rightarrow [0, 1]$ as follows. Set

$$k_0(\lambda) = \begin{cases} (1 - \frac{\varepsilon}{4})^{-1} \lambda & 0 \leq \lambda \leq 1 - \frac{\varepsilon}{4} \\ 1 & 1 - \frac{\varepsilon}{4} \leq \lambda \leq 1. \end{cases}$$

For $n = 1, 2, \dots, N$, set

$$k_n(\lambda) = \begin{cases} 0 & 0 \leq \lambda \leq 1 - 2^{-n-1}\varepsilon \\ 2^{n+2}\varepsilon^{-1}(\lambda - 1 + 2^{-n-1}\varepsilon) & 1 - 2^{-n-1}\varepsilon \leq \lambda \leq 1 - 2^{-n-2}\varepsilon \\ 1 & 1 - 2^{-n-2}\varepsilon \leq \lambda \leq 1. \end{cases}$$

Then $k_{n-1}k_n = k_n$ for $n = 1, 2, \dots, N$ and $|k_0(\lambda) - \lambda| \leq \frac{\varepsilon}{4}$ for all $\lambda \in [0, 1]$. For $n = 0, 1, \dots, N$, in Lemma 2.5 choose $\delta_n > 0$ for $f = k_n$ and with ε as given. Set

$$\delta = \min(\delta_0, \delta_1, \dots, \delta_N, \frac{\varepsilon}{2}).$$

Apply Definition 3.1 with δ in place of ε and with $m, a_1, a_2, \dots, a_m, x, y$ as given, getting $c_1, c_2, \dots, c_m \in A$ and $g \in B$ as there. For $n = 0, 1, \dots, N$, set $g_n = k_n(g)$. We verify the conclusions of the lemma.

Conclusion (1) is clear from the construction. Conclusion (2) is immediate. For conclusion (3), observe that $1 - g_N$ is gotten from $1 - g$ by functional calculus using the function $\lambda \mapsto 1 - k_N(1 - \lambda)$. This function vanishes at 0, so for $j = 1, 2, \dots, m$ we have $(1 - g_N)c_j \in B$ by Corollary 2.7. Now for $n = 0, 1, \dots, N - 1$, we have $(1 - g_n)c_j = (1 - g_n)(1 - g_N)c_j \in B$. For conclusion (4), use Lemma 2.2(i) of [12] to get $g_0 \preceq g$. For $n = 1, 2, \dots, N$, we then get $g_n \preceq g$ since $g_n \leq g_0$.

For conclusion (5), we start with the calculation

$$\|(1 - g_0)x(1 - g_0)\| \geq \|(1 - g)x(1 - g)\| - 2\|g - g_0\| > 1 - \frac{\varepsilon}{2} - 2\left(\frac{\varepsilon}{4}\right) = 1 - \varepsilon.$$

For $n = 1, 2, \dots, N$, use $(1 - g_n)^2 \geq (1 - g_0)^2$ at the second step to get

$$\begin{aligned} \|(1 - g_n)x(1 - g_n)\| &= \|x^{1/2}(1 - g_n)^2x^{1/2}\| \geq \|x^{1/2}(1 - g_0)^2x^{1/2}\| \\ &= \|(1 - g_0)x(1 - g_0)\| > 1 - \varepsilon. \end{aligned}$$

Conclusion (6) follows directly from the choice of δ_n . ■

Lemma 6.2. *Let $m, n, N \in \mathbb{Z}_{>0}$. Let A be an infinite dimensional simple separable unital C^* -algebra. Let $B \subset A$ be centrally large in A and suppose that $\text{tsr}(B) = 1$. Let $x_1, x_2, \dots, x_n \in A_+$ be positive elements of norm one. Let $a_1, a_2, \dots, a_m \in A$ be arbitrary. Then for all $\varepsilon > 0$ there exists $h \in B$ with $0 \leq h \leq 1$, there exist $r_{k,l} \in B$ for $k = 1, 2, \dots, n$ and for $l = 0, 1, \dots, N - 1$, and there exist $b_1, b_2, \dots, b_m \in A$ such that:*

- (1) $0 \leq r_{k,l} \leq 1$ and $\|r_{k,l}\| = 1$ for $k = 1, 2, \dots, n$ and $l = 0, 1, \dots, N$.
- (2) $\|x_k r_{k,0} - r_{k,0}\| < \varepsilon$ for $k = 1, 2, \dots, n$.
- (3) $1 - h \preceq_B r_{k,N}$ for $k = 1, 2, \dots, n$.

- (4) $r_{k,l}r_{k,l+1} = r_{k,l+1}$ for $k = 1, 2, \dots, n$ and $l = 0, 1, \dots, N-2$.
- (5) $hr_{k,0} = r_{k,0}$ for $k = 1, 2, \dots, n$.
- (6) $r_{k,l} \in \overline{hBh}$ for $k = 1, 2, \dots, n$ and $l = 0, 1, \dots, N$.
- (7) $\|a_j - b_j\| < \varepsilon$ for $j = 1, 2, \dots, m$.
- (8) $hb_j \in \overline{hB} \subset B$ for $j = 1, 2, \dots, m$.

Proof. The subalgebra B is simple by Proposition 5.2 of [16], finite by Proposition 3.1 of [22], and infinite dimensional by Proposition 5.5 of [16].

Set $\varepsilon_0 = \min(\frac{\varepsilon}{4}, \frac{1}{48})$. Define continuous functions $p_0, p_1: [0, 1] \rightarrow [0, 1]$ by

$$p_0(\lambda) = \begin{cases} (1 - 2\varepsilon_0)^{-1}\lambda & 0 \leq \lambda \leq 1 - 2\varepsilon_0 \\ 1 & 1 - 2\varepsilon_0 \leq \lambda \leq 1 \end{cases}$$

and

$$p_1(\lambda) = \begin{cases} 0 & 0 \leq \lambda \leq 1 - 2\varepsilon_0 \\ \varepsilon_0^{-1}(\lambda - 1 + 2\varepsilon_0) & 1 - 2\varepsilon_0 \leq \lambda \leq 1 - \varepsilon_0 \\ 1 & 1 - \varepsilon_0 \leq \lambda \leq 1. \end{cases}$$

Define $x_k^{(0)} = p_0(x_k)$ and $x_k^{(1)} = p_1(x_k)$ for $k = 1, 2, \dots, n$. This gives

$$(6.1) \quad \|x_k - x_k^{(0)}\| \leq 2\varepsilon_0, \quad x_k^{(0)}x_k^{(1)} = x_k^{(1)}, \quad \text{and} \quad \|x_k^{(1)}\| = 1.$$

We further need the functions $f_{1/2^{N+2}}, f_{1/2^{N+1}}, \dots, f_{1/4}: [0, 1] \rightarrow [0, 1]$ of Definition 2.3. We name them instead, in order, $f_0, f_1, \dots, f_N$, getting

$$(6.2) \quad f_l f_{l+1} = f_{l+1}$$

for $l = 0, 1, \dots, N-1$ and

$$(6.3) \quad f_N|_{[\frac{1}{2}, 1]} = 1.$$

For $k = 1, 2, \dots, n$, use Lemma 5.3 of [16] (with $r = 1$) to choose $y_k^{(0)} \in B_+$ such that

$$\|y_k^{(0)}\| = 1 \quad \text{and} \quad \|(x_k^{(1)})^{1/2} y_k^{(0)} - y_k^{(0)}\| < \frac{1}{48}.$$

Define

$$y_k = p_0(y_k^{(0)}) \quad \text{and} \quad z_k = p_1(y_k^{(0)}).$$

Then $y_k z_k = z_k$ and $\|z_k\| = 1$. Also,

$$\|y_k - y_k^{(0)}\| \leq 2\varepsilon_0 \leq \frac{1}{24},$$

so

$$(6.4) \quad \|(x_k^{(1)})^{1/2} y_k - y_k\| \leq 2\|y_k - y_k^{(0)}\| + \|(x_k^{(1)})^{1/2} y_k^{(0)} - y_k^{(0)}\| < \frac{1}{12} + \frac{1}{48} = \frac{5}{48}.$$

For $k = 1, 2, \dots, n$, Lemma 2.2 or Lemma 2.3 of [16] (depending on whether $z_k B z_k$ is unital) provides nonzero orthogonal unitarily equivalent positive elements $c_{k,1}, c_{k,2} \in \overline{z_k B z_k}$. We may assume that $\|c_{k,1}\| = \|c_{k,2}\| = 1$. Since B is finite and infinite dimensional, we can use Lemma 2.9 of [16] to find $d_k \in B_+ \setminus \{0\}$ such that whenever $g \in B$ satisfies $0 \leq g \leq 1$ and $g \prec_B d_k$, then

$$(6.5) \quad \|(1 - g)z_k(1 - g)\| > \frac{5}{6}.$$

Now use simplicity of B and Lemma 2.6 of [16] to find a nonzero positive element $c \in B$ such that for $k = 1, 2, \dots, n$ we have

$$(6.6) \quad c \preceq_B d_k \quad \text{and} \quad c \preceq_B \left(c_{k,1} - \frac{3}{4}\right)_+.$$

By a polynomial approximation argument, there is $\delta_0 > 0$ such that whenever D is a C^* -algebra and $y_1, y_2 \in D$ satisfy $0 \leq y_1, y_2 \leq 1$ and $\|y_1 - y_2\| < 2\delta_0$, then $\|f_0(y_1) - f_0(y_2)\| < \frac{\varepsilon}{4}$. Set $\delta = \min(\varepsilon, \delta_0, \frac{1}{48})$. Apply Lemma 6.1 with $N = 3$ and δ in place of ε , getting $g_0, g_1, g_2, g_3 \in B$ and $b_1, b_2, \dots, b_m \in A$ such that:

- (9) $0 \leq g_0, g_1, g_2, g_3 \leq 1$, $g_0 g_1 = g_1$, $g_1 g_2 = g_2$, and $g_2 g_3 = g_3$.
- (10) $\|b_j - a_j\| < \delta$ and $(1 - g_3)b_j \in B$ for $j = 1, 2, \dots, m$.
- (11) $g_0 \preceq_B c$.
- (12) $\|[g_1, (x_k^{(1)})^{1/2}]\| < \delta$ and $\|[g_2, (x_k^{(1)})^{1/2}]\| < \delta$ for $k = 1, 2, \dots, n$.

Part (7) of the conclusion is immediate from the choice of $b_1, b_2, \dots, b_m$. Set $h = 1 - g_3$. Then $hb_j \in B$ by (10). Thus $\lim_{n \rightarrow \infty} h^{1/n} h b_j = h b_j \in \overline{hB}$, which is part (8) of the conclusion.

For $k = 1, 2, \dots, n$, further define $r_k = (1 - g_2)x_k^{(1)}(1 - g_2)$ and, for $l = 0, 1, \dots, N$, define $r_{k,l} = f_l(r_k)$. Clearly $0 \leq r_{k,l} \leq 1$. From (6.2) we get

$$r_{k,l} r_{k,l+1} = r_{k,l+1}$$

for $l = 0, 1, \dots, N - 1$. This is part (4) and most of part (1) of the conclusion. We next use (6.4) twice to get

$$(6.7) \quad \|(x_k^{(1)})^{1/2} y_k (x_k^{(1)})^{1/2} - y_k\| < \frac{5}{24}.$$

Using this relation at the second step, $z_k \leq y_k$ at the third step, and $g_2 \leq g_0$, (11), (6.6), and the choice of d_k (which gave (6.5)) at the fourth step, we get the estimate

$$\begin{aligned} \|r_k\| &\geq \|(1 - g_2)(x_k^{(1)})^{1/2} y_k (x_k^{(1)})^{1/2} (1 - g_2)\| > \|(1 - g_2)y_k(1 - g_2)\| - \frac{5}{24} \\ &\geq \|(1 - g_2)z_k(1 - g_2)\| - \frac{5}{24} > \frac{5}{6} - \frac{5}{24} > \frac{1}{2}. \end{aligned}$$

In particular, by (6.3), we have $\|r_{k,N}\| = 1$. This implies that $\|r_{k,l}\| \geq 1$ for $r_{k,l} r_{k,l+1} = r_{k,l+1}$, and finishes the proof of part (1) of the conclusion.

Part (5) of the conclusion follows from $h = 1 - g_3$, $g_2 g_3 = g_3$, and the fact that $r_{k,0} \in \overline{(1 - g_2)B(1 - g_2)}$. Combining this with part (4), we get part (6).

Our next goal is part (3) of the conclusion. We start with

$$(6.8) \quad 1 - h = g_3 \leq g_0.$$

We next claim that

$$(6.9) \quad g_0 \preceq_B \left[(1 - g_1)y_k(1 - g_1) - \frac{3}{4}\right]_+.$$

We use (11) and (6.6) at the first step, Lemma 1.7 of [16], $c_{k,1}c_{k,2} = 0$, unitary equivalence of $c_{k,1}$ and $c_{k,2}$, and $c_{k,1} + c_{k,2} \leq y_k$ at the second step, Lemma 1.8 of [16] at the third step, and $g_0 g_1 = g_1$ and Lemma 1.10 of [16] at the fourth step,

getting

$$\begin{aligned}
g_0 \oplus g_0 &\preceq_B (c_{k,1} - \frac{3}{4})_+ + (c_{k,2} - \frac{3}{4})_+ \\
&\preceq_B (y_k - \frac{3}{4})_+ \\
&\preceq_B [(1 - g_1)y_k(1 - g_1) - \frac{3}{4}]_+ \oplus g_1 \\
&\preceq_B [(1 - g_1)y_k(1 - g_1) - \frac{3}{4}]_+ \oplus (g_0 - \frac{1}{2})_+.
\end{aligned}$$

Since $\text{tsr}(B) = 1$, we can apply Theorem 4.3 of [26] to get (6.9), as desired.

Now we claim that

$$(6.10) \quad [(1 - g_1)y_k(1 - g_1) - \frac{3}{4}]_+ \preceq_A (r_k - \frac{1}{2})_+.$$

Set

$$s = [(1 - g_2)(x_k^{(1)})^{1/2}(1 - g_2)][(1 - g_1)y_k(1 - g_1)][(1 - g_2)(x_k^{(1)})^{1/2}(1 - g_2)].$$

Then, using $(1 - g_1)(1 - g_2) = 1 - g_1$ at the first step, $\|1 - g_2\| \leq 1$ and (12) at the third step, (6.7) and $\|1 - g_1\| \leq 1$ at the fifth step, and the choice of δ at the sixth step,

$$\begin{aligned}
&\|s - (1 - g_1)y_k(1 - g_1)\| \\
&= \|(1 - g_2)(x_k^{(1)})^{1/2}(1 - g_1)y_k(1 - g_1)(x_k^{(1)})^{1/2}(1 - g_2) \\
&\quad - (1 - g_2)(1 - g_1)y_k(1 - g_1)(1 - g_2)\| \\
&\leq \|1 - g_2\|^2 \|(x_k^{(1)})^{1/2}(1 - g_1)y_k(1 - g_1)(x_k^{(1)})^{1/2} - (1 - g_1)y_k(1 - g_1)\| \\
&< 2\delta + \|(1 - g_1)(x_k^{(1)})^{1/2}y_k(x_k^{(1)})^{1/2}(1 - g_1) - (1 - g_1)y_k(1 - g_1)\| \\
&\leq 2\delta + \|1 - g_1\|^2 \|(x_k^{(1)})^{1/2}y_k(x_k^{(1)})^{1/2} - y_k\| \\
&< 2\delta + \frac{5}{24} \leq \frac{1}{4}.
\end{aligned}$$

So Corollary 1.6 of [16] implies that

$$[(1 - g_1)y_k(1 - g_1) - \frac{3}{4}]_+ \preceq_A (s - \frac{1}{2})_+.$$

Clearly

$$s \leq (1 - g_2)(x_k^{(1)})^{1/2}(x_k^{(1)})^{1/2}(1 - g_2) = r_k,$$

so

$$(s - \frac{1}{2})_+ \preceq_A (r_k - \frac{1}{2})_+$$

by Lemma 1.7 of [16]. The claim follows.

Using (6.3) and the definition $r_{k,N} = f_N(r_k)$, we get

$$r_{k,N}(r_k - \frac{1}{2})_+ = (r_k - \frac{1}{2})_+.$$

In particular, $(r_k - \frac{1}{2})_+ \preceq_A r_{k,N}$. Combining this relation with (6.8), (6.9), and (6.10), we get $1 - h \preceq_A r_{k,N}$. This is part (3) of the conclusion.

It remain to prove part (2) of the conclusion. It follows from (12) that

$$\|(x_k^{(1)})^{1/2}(1 - g_2)^2(x_k^{(1)})^{1/2} - (1 - g_2)x_k^{(1)}(1 - g_2)\| < 2\delta.$$

Using $\delta \leq \delta_0$, we get

$$(6.11) \quad \|f_0((x_k^{(1)})^{1/2}(1 - g_2)^2(x_k^{(1)})^{1/2}) - f_0((1 - g_2)x_k^{(1)}(1 - g_2))\| < \frac{\varepsilon}{4}.$$

Since $x_k^{(0)}x_k^{(1)} = x_k^{(1)}$ (from (6.1)), we get

$$x_k^{(0)}f_0((x_k^{(1)})^{1/2}(1-g_2)^2(x_k^{(1)})^{1/2}) = f_0((x_k^{(1)})^{1/2}(1-g_2)^2(x_k^{(1)})^{1/2}).$$

Combining this equation with $\|x_k - x_k^{(0)}\| \leq 2\varepsilon_0$ (also from (6.1)) and using (6.11) twice, we now get

$$\|x_k f_0((1-g_2)x_k^{(1)}(1-g_2)) - f_0((1-g_2)x_k^{(1)}(1-g_2))\| < 2\varepsilon_0 + \frac{2\varepsilon}{4} \leq \varepsilon.$$

By the definition of $r_{k,0}$, we thus have $\|x_k r_{k,0} - r_{k,0}\| < \varepsilon$, as desired. $\blacksquare$

Theorem 6.3. *Suppose A is an infinite dimensional simple separable unital C^* -algebra. Suppose A has a centrally large subalgebra B such that $\text{tsr}(B) = 1$. Then $\text{tsr}(A) = 1$.*

Proof. We will show that every two sided zero divisor in A is a limit of invertible elements. That is, if $a \in A$ and there are nonzero $x, y \in A$ such that $xa = ay = 0$, then we show that for every $\varepsilon > 0$ there is an invertible element $z \in A$ such that $\|z - a\| < \varepsilon$. It will follow from Theorem 3.3(a) of [23] (see Definition 3.1 of [23]) that any $a \in A$ which is not a limit of invertible elements is left or right invertible but not both. Now B has stable rank one, so is finite by Proposition 3.1 of [22]. Proposition 6.15 of [16] then implies that A is finite, so there are no such elements. Thus we conclude that A has stable rank one.

Without loss of generality, $\|a\| \leq 1/2$ and $\varepsilon \leq 1$. Replacing x with $\|x\|^{-2}x^*x$ and y with $\|y\|^{-2}yy^*$, we may assume that x and y are positive elements of norm 1.

Set $\delta_1 = \min(1, \varepsilon)/20$. Apply Lemma 6.2 with $N = 1$, with $n = 2$, $x_1 = x$, and $x_2 = y$, with $m = 1$ and $a_1 = a$, and with δ_1 in place of ε , to get $a_0 \in A$, $h_1 \in B$ with $0 \leq h_1 \leq 1$, and $s_0, s_1, t_0, t_1 \in \overline{h_1 B h_1}$ such that:

- (1) $0 \leq s_0, s_1, t_0, t_1 \leq 1$ and $\|s_0\| = \|s_1\| = \|t_0\| = \|t_1\| = 1$.
- (2) $\|xs_0 - s_0\| < \delta_1$ and $\|yt_0 - t_0\| < \delta_1$.
- (3) $1 - h_1 \preceq_B s_1$ and $1 - h_1 \preceq_B t_1$.
- (4) $s_0 s_1 = s_1$ and $t_0 t_1 = t_1$.
- (5) $h_1 s_0 = s_0$ and $h_1 t_0 = t_0$.
- (6) $\|a - a_0\| < \delta_1$.
- (7) $h_1 a_0 \in \overline{h_1 B} \subset B$.

Set $a_1 = (1 - s_0)a_0(1 - t_0)$. Then $s_1 a_1 = (s_1 - s_1 s_0)a_0(1 - t_0) = 0$ by (4) and selfadjointness, and similarly $a_1 t_1 = 0$.

Next we estimate $\|a_1 - a\|$. Since $\|a\| \leq \frac{1}{2}$ and $xa = 0$, we have

$$\|s_0 a_0\| \leq \|s_0\| \cdot \|a_0 - a\| + \|s_0 - s_0 x\| \cdot \|a\| \leq 2\delta_1.$$

Similarly $\|a_0 t_0\| < 2\delta_1$. Now we can compute

$$\begin{aligned} \|a - a_1\| &\leq \|a - a_0\| + \|a_0 - (1 - s_0)a_0(1 - t_0)\| \\ &< \delta_1 + \|a_0 t_0\| + \|s_0 a_0\| \cdot \|1 - t_0\| \leq 5\delta_1. \end{aligned}$$

This also gives us $\|a_1\| < 5\delta_1 + \frac{1}{2} \leq 1$.

The subalgebra B is simple by Proposition 5.2 of [16]. Apply Lemma 2.4, getting $r \in B_+ \setminus \{0\}$ with $r \leq s_1$ and $r \preceq_B t_1$. Set $\delta_2 = \frac{1}{2}\|r\|$, let f_{δ_2} be as in Definition 2.3, and set $q = f_{\delta_2}(r)$. Thus $\|q\| = 1$. The functions f_ε of Definition 2.3 are the same as those defined at the beginning of Section 2 of [24]. Since $\text{tsr}(B) = 1$, Proposition 2.4(v) in [24] therefore provides a unitary $v \in B$ such that $v^* q v \in \overline{t_1 B t_1}$.

We claim that

$$(6.12) \quad qa_1v^* = a_1v^*q = 0.$$

For the first, use $q \in \overline{s_1Bs_1}$ and $s_1a_1 = 0$ to get $qa_1 = 0$. For the second, use $v^*qv \in \overline{t_1Bt_1}$ and $a_1t_1 = 0$ to get $a_1v^*qv = 0$, and multiply on the right by v^* to get $a_1v^*q = 0$. The claim is proved.

Set

$$\delta_3 = \min \left(\frac{\varepsilon}{20}, \frac{1}{4} \right).$$

Apply Lemma 6.2 with $N = 1$, with $n = 1$ and $x_1 = q$, with $m = 1$ and a_1v^* in place of $a_1, a_2, \dots, a_m$, and with δ_3 in place of ε , to get $a_2 \in A$, $h_2 \in B$ with $0 \leq h_2 \leq 1$, and $r_0, r_1 \in \overline{h_2Bh_2}$ such that:

- (8) $0 \leq r_0, r_1 \leq 1$ and $\|r_0\| = \|r_1\| = 1$.
- (9) $\|qr_0 - r_0\| < \delta_3$.
- (10) $1 - h_2 \preceq_B r_1$.
- (11) $r_0r_1 = r_1$.
- (12) $h_2r_0 = r_0$.
- (13) $\|a_2 - a_1v^*\| < \delta_3$.
- (14) $h_2a_2 \in \overline{h_2B} \subset B$.

Define $a_3 = (1 - r_0)a_2(1 - r_0)$. Then $r_1a_3 = a_3r_1 = 0$. Next,

$$\|a_2\| \leq \|a_2 - a_1v^*\| + \|a_1 - a\| \cdot \|v^*\| + \|a\| \cdot \|v^*\| < \delta_3 + 5\delta_1 + 1/2 \leq 1.$$

From (9) and selfadjointness, we get $\|r_0 - r_0q\| < \delta_3$. Using this and $qa_1v^* = 0$ (from (6.12)) at the second step, we have

$$\begin{aligned} \|r_0a_2\| &\leq \|r_0a_2 - r_0a_1v^*\| + \|r_0 - r_0q\| \cdot \|a_1v^*\| + \|r_0qa_1v^*\| \\ &\leq \delta_3\|r_0\| + \delta_3\|a_1\| \cdot \|v^*\| \leq 2\delta_3. \end{aligned}$$

Similarly, $a_1v^*q = 0$ (also from (6.12)) gives $\|a_2r_0\| \leq 2\delta_3$.

Now we can estimate:

$$\|a_2 - a_3\| = \|a_2 - (1 - r_0)a_2(1 - r_0)\| \leq \|a_2r_0\| + \|r_0a_2\| \cdot \|1 - r_0\| < 4\delta_3.$$

Therefore

$$\|a_3v - a\| \leq \|a_3 - a_2\| \cdot \|v\| + \|a_2v - a\| + \|a_1 - a\| < 4\delta_3 + \delta_3 + 5\delta_1 \leq \frac{\varepsilon}{2}.$$

Combining (12) with its adjoint, we get $h_2r_0 = r_0h_2$. So $h_2^{1/2}r_0 = r_0h_2^{1/2}$.

Therefore

$$h_2a_3 = h_2^{1/2}(1 - r_0)h_2^{1/2}a_2(1 - r_0).$$

We have $h_2^{1/2}a_2 \in B$ by (14) and Corollary 2.7, and $1 - r_0 \in B$, so

$$(6.13) \quad h_2a_3 \in \overline{h_2B} \cdot \overline{h_2B} \cdot B \subset \overline{h_2B} \subset B.$$

If $h_2 = 1$, then $a_3 \in B$. Since $\text{tsr}(B) = 1$, there is an invertible element $z_0 \in B$ such that $\|z_0 - a_3\| < \varepsilon - \|a_3v - a\|$. Set $z = z_0v$. Then z is invertible in A and $\|z - a\| < \varepsilon$, as desired. So we may assume that $h_2 \neq 1$. Choose $\delta_4 > 0$ such that $\text{sp}(h_2)$ is not contained in $[1 - 2\delta_4, 1]$. In particular, $\delta_4 < \frac{1}{2}$. Let f_{δ_4} be as in Definition 2.3. Since $1 - h_2 \geq 0$ and $\text{sp}(1 - h_2)$ is not contained in $[0, 2\delta_4]$, we have $\|f_{\delta_4}(1 - h_2)\| = 1$.

Using (10), $\text{tsr}(B) = 1$, and Proposition 2.4(v) in [24], we get a unitary $u \in U(B)$ such that

$$(6.14) \quad uf_{\delta_4}(1 - h_2)u^* \in \overline{r_1 B r_1}.$$

Define

$$z_1 = uf_{\delta_4}(1 - h_2)u^*, \quad z_3 = f_{\delta_4}(1 - h_2), \quad \text{and} \quad z_2 = 1 - z_1 - z_3.$$

We want to apply Lemma 5.4 with A , B , z_1 , z_2 , and z_3 as given, and with $a_3 u$ in place of a .

We have $\text{tsr}(B) = 1$ by hypothesis, and hypothesis (2) of Lemma 5.4 holds by definition. Also $z_1, z_2, z_3 \in B$ by construction.

Next, we claim that $z_1 z_3 = 0$. From (11) and (12), we get $h_2 r_1 = r_1$, so $(1 - h_2)r_1 = 0$. Thus $z_1(1 - h_2) = uf_{\delta_4}(1 - h_2)u^*(1 - h_2) = 0$ by Lemma 5.1, and another application of Lemma 5.1 gives

$$(6.15) \quad z_1 z_3 = z_1 f_{\delta_4}(1 - h_2)z_1 = 0,$$

proving the claim. This is hypothesis (3) of Lemma 5.4. Clearly $0 \leq z_1 \leq 1$ and $0 \leq z_3 \leq 1$, so we get $z_2 \geq 0$. Thus hypothesis (1) of Lemma 5.4 holds.

Combining (6.14) with (11) and the definition of z_1 , we get $r_0 z_1 = z_1 r_0 = z_1$. So

$$z_1 a_3 u = z_1(1 - r_0)a_2(1 - r_0)u = 0,$$

which is hypothesis (4) of Lemma 5.4. Clearly $u z_3 = z_1 u$, so also

$$a_3 u z_3 = a_3 z_1 u = (1 - r_0)a_2(1 - r_0)z_1 u = 0,$$

which is hypothesis (5) of Lemma 5.4.

It remains only to verify hypothesis (6) of Lemma 5.4. Recall from (6.13) that $h_2 a_3 \in B$. Since $\delta_4 < \frac{1}{2}$, the function $\lambda \mapsto 1 - f_{\delta_4}(1 - \lambda)$, from $[0, 1]$ to $[0, 1]$, vanishes at zero. So Corollary 2.7 implies $(1 - z_3)a_3 = [1 - f_{\delta_4}(1 - h_2)]a_3 \in B$. From (6.15) and $z_1, u, z_2 \in B$, we now get $z_2 a_3 u z_2 = (1 - z_1)(1 - z_3)a_3 u z_2 \in B$, as desired.

We can now apply Lemma 5.4 to find an invertible element $a_4 \in A$ such that $\|a_4 - a_3 u\| < \frac{\varepsilon}{2}$. Then $z = a_4 u^* v$ is invertible and satisfies

$$\|z - a\| \leq \|a_4 - a_3 u\| \cdot \|u^*\| \cdot \|v\| + \|a_3 v - a\| < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon.$$

This completes the proof. $\blacksquare$

Combining Theorem 6.3 with results of [3] and [16], we can show that if, in addition, the centrally large subalgebra has real rank zero, then so does the containing algebra. It is possible that one does not need the subalgebra to have stable rank one, but the proof would need to be much longer.

Theorem 6.4. *Suppose A is an infinite dimensional simple separable unital C^* -algebra. Suppose A has a centrally large subalgebra B such that $\text{tsr}(B) = 1$ and $\text{RR}(B) = 0$. Then $\text{RR}(A) = 0$.*

Proof. We use the Cuntz semigroups $\text{Cu}(B)$ and $\text{Cu}(A)$. We refer to [2] and Section 1 of [16] for background. In particular, recall the compact containment relation $\ll$, Definition 1.24 and Lemma 1.25 of [16] or (under the name “way below”) condition **(O4)** in Definition 4.1 of [2]. If D is a C^* -algebra, we further call an element $\eta \in \text{Cu}(D)$ compact if $\eta \ll \eta$.

Since $\text{RR}(B) = 0$ and $\text{tsr}(B) = 1$, it follows from the second paragraph of Corollary 5 on page 186 of [3] that every element of $\text{Cu}(B)$ is the supremum of a nondecreasing sequence of compact elements. We claim that the same is true of A . So let $\eta \in \text{Cu}(A)$. If η is the class of a projection in $K \otimes A$, then η is itself compact, by Lemma 1.25(3) of [16]. So the claim is immediate. Otherwise, let $\iota: B \rightarrow A$ be the inclusion map. It follows from Theorem 6.3 that $\text{tsr}(A) = 1$, so A is stably finite by Proposition 3.1 and Theorem 3.3 of [22]. Since centrally large algebras are large, Corollary 3.6 now implies that B is stably large in A . So Theorem 6.6 of [16] provides $\mu \in \text{Cu}(B)$ such that $\iota_*(\mu) = \eta$. Let $(\mu_n)_{n \in \mathbb{Z}_{>0}}$ be a nondecreasing sequence of compact elements in $\text{Cu}(B)$ such that $\mu = \sup_{n \in \mathbb{Z}_{>0}} \mu_n$. Then $(\iota_*(\mu_n))_{n \in \mathbb{Z}_{>0}}$ is a nondecreasing sequence of compact elements in $\text{Cu}(A)$ such that $\eta = \sup_{n \in \mathbb{Z}_{>0}} \iota_*(\mu_n)$. This proves the claim.

We saw already that $\text{tsr}(A) = 1$. Combining this fact, the claim, and Corollary 5 on page 186 of [3], we get $\text{RR}(A) = 0$. ■

7. APPLICATION TO THE STABLE RANK OF CROSSED PRODUCTS

In this section, we apply our main result to show that the crossed product by a minimal homeomorphism of a compact metric space X has stable rank one whenever there is a continuous surjective map from X to the Cantor set. We conjecture that the result holds whenever X is infinite. Our theorem covers examples in which it is known that the crossed product is not Z -stable, and in particular is not classifiable in terms of K -theory and traces. In particular, our result can't be proved by using the fact that simple unital Z -stable C^* -algebras have stable rank one (Theorem 6.7 of [25]).

Theorem 7.1. *Let X be a compact metric space. Assume that there is a continuous surjective map from X to the Cantor set. Let $h: X \rightarrow X$ be a minimal homeomorphism. Then $C^*(\mathbb{Z}, X, h)$ has stable rank one.*

In Proposition 5.1 of [11], it is shown that the conditions on the pair (X, h) are equivalent to a number of other conditions. For example, the hypotheses are equivalent to the assumption that (X, h) has a factor system which is a homeomorphism of the Cantor set. They are also equivalent to the assumption that there is a decreasing sequence $Y_0 \supset Y_1 \supset \cdots$ of nonempty compact open subsets of X such that the subset $Y = \bigcap_{n=0}^{\infty} Y_n$ satisfies $h^r(Y) \cap Y = \emptyset$ for all $r \in \mathbb{Z} \setminus \{0\}$.

The proof depends on a result in [7], according to which simple direct limits of AH systems with diagonal maps have stable rank one, regardless of any conditions on dimension growth. It seems plausible to believe that the same should be true for simple direct limits of recursive subhomogeneous algebras in which the maps of the system are diagonal. For any minimal homeomorphism h of an infinite compact metric space, and any nonempty closed subset $Y \subset X$ such that $h^r(Y) \cap Y = \emptyset$ for all $r \in \mathbb{Z} \setminus \{0\}$, the centrally large subalgebra $C^*(\mathbb{Z}, X, h)_Y$ (see the introduction for the notation) is the direct limit of such a system. (This will appear in [19].) Accordingly, we make the following conjecture.

Conjecture 7.2 (joint with Zhuang Niu). *Let X be an infinite compact metric space and let $h: X \rightarrow X$ be a minimal homeomorphism. Then $C^*(\mathbb{Z}, X, h)$ always has stable rank one.*

Since simple unital AH algebras can have arbitrary stable rank (even infinite stable rank; see Theorems 8 and 12 of [28]), this would show that some kinds of

pathology which occur for simple unital AH algebras are not possible with crossed products by minimal homeomorphisms.

Proof of Theorem 7.1. Proposition 5.14 of [11] provides a nonempty compact subset $Y \subset X$ such that $h^n(Y) \cap Y = \emptyset$ for all $n \in \mathbb{Z} \setminus \{0\}$ and such that $C^*(\mathbb{Z}, X, h)_Y$ (as defined in the introduction) is the direct limit of an AH system with diagonal maps in the sense described at the beginning of Section 2.2 of [7]. (The maps are explicitly described in Lemma 5.12 of [11].) By Theorem 7.10 of [16], the subalgebra $C^*(\mathbb{Z}, X, h)_Y$ is a large subalgebra of $C^*(\mathbb{Z}, X, h)$ of crossed product type in the sense of Definition 4.9 of [16]. In particular, $C^*(\mathbb{Z}, X, h)_Y$ is simple (by Proposition 5.2 of [16]) and is centrally large in $C^*(\mathbb{Z}, X, h)$ (by Theorem 4.6). Using simplicity and the diagonal maps condition in the AH system, it follows from Theorem 4.1 of [7] that $C^*(\mathbb{Z}, X, h)_Y$ has stable rank one. So $C^*(\mathbb{Z}, X, h)$ has stable rank one by Theorem 6.3. ■

Corollary 7.3. *Let Y be a compact metric space with more than one point, let $d \in (0, 1)$, and let X and the minimal homeomorphism $h: X \rightarrow X$ be a subshift (of the shift on $(Y \times [0, 1])^{\mathbb{Z}}$) as constructed in Section 2 of [10]. Then $C^*(\mathbb{Z}, X, h)$ has stable rank one.*

Proof. On the bottom of page 111 of [10], it is proved that the system (X, h) has the universal odometer as a factor system. (The hypothesis that Y be contractible, which is being assumed at this point, is not needed for this conclusion.) In particular, there is a continuous surjective map from X to the Cantor set. So Theorem 7.1 applies. ■

Example 7.4. *Let $n \in \mathbb{Z}_{>0}$ satisfy $n \geq 2$. In Corollary 7.3, make the particular choices $d \in (1 - \frac{1}{n}, 1)$ and $Y = [0, 1]^{3n}$. Theorem 2.2 of [10] implies that the radius of comparison of $C^*(\mathbb{Z}, X, h)$ satisfies $\text{rc}(C^*(\mathbb{Z}, X, h)) \geq n - 1$. In particular, $C^*(\mathbb{Z}, X, h)$ does not have strict comparison, so is not Z -stable (by Corollary 4.6 of [25]), but has stable rank one by Corollary 7.3.*

The mean dimension $\text{mdim}(h)$ of a homeomorphism h of a compact metric space is defined in Definition 2.6 of [14]. It is shown in [11] that for the minimal homeomorphisms in Corollary 7.3 one actually has $\text{rc}(C^*(\mathbb{Z}, X, h)) = \frac{1}{2} \text{mdim}(h)$. It is pointed out after Proposition 2.4 of [10] that if the covering dimension of the space Y in Corollary 7.3 satisfies $\dim(Y) > 0$, then the homeomorphism h of Corollary 7.3 satisfies $\text{mdim}(h) > 0$. Thus, in fact, whenever $\dim(Y) > 0$ in Corollary 7.3, then $\text{rc}(C^*(\mathbb{Z}, X, h)) > 0$, and $C^*(\mathbb{Z}, X, h)$ is not Z -stable (but does have stable rank one).

Example 7.5. *In Corollary 7.3, make the particular choices $d \in (0, 1)$ and $Y = S^2 \times S^2$. Theorem 3.1 of [10] implies that $K_0(C^*(\mathbb{Z}, X, h))$ is not weakly unperforated. We can now give a different argument to show that $C^*(\mathbb{Z}, X, h)$ is not Z -stable. By Corollary 4.9 of [25], if $C^*(\mathbb{Z}, X, h)$ were Z -stable then (using simplicity of $C^*(\mathbb{Z}, X, h)$ to get simplicity of $V(C^*(\mathbb{Z}, X, h))$) it would follow that $K_0(C^*(\mathbb{Z}, X, h))$ is almost unperforated. By the discussion after Lemma 3.4 of [25], it would follow that $K_0(C^*(\mathbb{Z}, X, h))$ is weakly unperforated, a contradiction. Again, $C^*(\mathbb{Z}, X, h)$ has stable rank one by Corollary 7.3.*

There are other ways to construct examples of minimal homeomorphisms for which we can prove that the mean dimension is nonzero and that the crossed

products have stable rank one, although we do not know that the crossed products don't have strict comparison, and we do not know that they are not Z -stable.

Example 7.6. *Let X_0 be any connected compact metric space, and let $h_0: X_0 \rightarrow X_0$ be a minimal homeomorphism such that $\text{mdim}(h_0) > 0$. Let $k: Z \rightarrow Z$ be an odometer homeomorphism of the Cantor set. (See, for example, Section VIII.4 of [6].) Set $X = Z \times X_0$ and define $h = k \times h_0: X \rightarrow X$. It is shown in [11] that h is minimal. It is also shown in [11] that $\text{mdim}(h) = \text{mdim}(h_0)$. Obviously X has a continuous surjective map to the Cantor set, so $C^*(\mathbb{Z}, X, h)$ has stable rank one by Theorem 6.3.*

REFERENCES

- [1] D. Archey and N. C. Phillips, *Centrally large subalgebras and Z -stability*, in preparation.
- [2] P. Ara, F. Perera, and A. S. Toms, *K-Theory for operator algebras. Classification of C^* -algebras*, pages 1–71 in: *Aspects of Operator Algebras and Applications*, P. Ara, F. Lledó, and F. Perera (eds.), Contemporary Mathematics vol. 534, Amer. Math. Soc., Providence RI, 2011.
- [3] K. T. Coward, G. A. Elliott, and C. Ivanescu, *The Cuntz semigroup as an invariant for C^* -algebras*, J. reine angew. Math. **623**(2008), 161–193.
- [4] J. Cuntz, *The structure of multiplication and addition in simple C^* -algebras*, Math. Scand. **40**(1977), 215–233.
- [5] J. Cuntz, *Dimension functions on simple C^* -algebras*, Math. Ann. **233**(1978), 145–153.
- [6] K. R. Davidson, *C^* -Algebras by Example*, Fields Institute Monographs no. 6, Amer. Math. Soc., Providence RI, 1996.
- [7] G. A. Elliott, T. M. Ho, and A. S. Toms, *A class of simple C^* -algebras with stable rank one*, J. Funct. Anal. **256**(2009), 307–322.
- [8] G. A. Elliott and Z. Niu, *All irrational extended rotation algebras are AF algebras*, preprint.
- [9] G. A. Elliott and Z. Niu, *C^* -algebra of a minimal homeomorphism of zero mean dimension*, preprint (arXiv:1406.2382v2 [math.OA]).
- [10] J. Giol and D. Kerr, *Subshifts and perforation*, J. reine angew. Math. **639**(2010), 107–119.
- [11] T. Hines, N. C. Phillips, and A. S. Toms, *Mean dimension and radius of comparison for minimal homeomorphisms with Cantor factors*, in preparation.
- [12] E. Kirchberg and M. Rørdam, *Non-simple purely infinite C^* -algebras*, Amer. J. Math. **122**(2000), 637–666.
- [13] H. Lin, *An Introduction to the Classification of Amenable C^* -Algebras*, World Scientific, River Edge NJ, 2001.
- [14] E. Lindenstrauss and B. Weiss, *Mean topological dimension*, Israel J. Math. **115**(2000), 1–24.
- [15] N. C. Phillips, *Crossed products of the Cantor set by free minimal actions of $\mathbb{Z}^d$* , Commun. Math. Phys. **256**(2005), 1–42.
- [16] N. C. Phillips, *Large subalgebras*, preprint (arXiv: 1408.5546v1 [math.OA]).
- [17] N. C. Phillips, *The transformation group C^* -algebras of free minimal actions of $\mathbb{Z}^d$ on finite dimensional compact metric spaces (draft)*.
- [18] N. C. Phillips, *Stable rank one for free minimal actions of $\mathbb{Z}^d$ with Cantor factors*, in preparation.
- [19] N. C. Phillips, in preparation.
- [20] I. F. Putnam, *The C^* -algebras associated with minimal homeomorphisms of the Cantor set*, Pacific J. Math. **136**(1989), 329–353.
- [21] I. F. Putnam, *On the K-theory of C^* -algebras of principal groupoids*, Rocky Mountain J. Math. **28**(1998), 1483–1518.
- [22] M. A. Rieffel, *Dimension and stable rank in the K-theory of C^* -algebras*, Proc. London Math. Soc. (3) **46**(1983), 301–333.
- [23] M. Rørdam, *On the structure of simple C^* -algebras tensored with a UHF-algebra*, J. Funct. Anal. **100**(1991), 1–17.
- [24] M. Rørdam, *On the structure of simple C^* -algebras tensored with a UHF-algebra. II*, J. Funct. Anal. **107**(1992), 255–269.

- [25] M. Rørdam, *The stable and the real rank of $\mathcal{Z}$ -absorbing C^* -algebras*, International J. Math. **15**(2004), 1065–1084.
- [26] M. Rørdam and W. Winter, *The Jiang-Su algebra revisited*, J. reine angew. Math. **642**(2010), 129–155.
- [27] G. Szabó, *The Rokhlin dimension of topological $\mathbb{Z}^m$ -actions*, preprint (arXiv: 1308.5418v4 [math.OA]).
- [28] J. Villadsen, *On the stable rank of simple C^* -algebras*, J. Amer. Math. Soc. **12**(1999), 1091–1102.

DEPARTMENT OF MATHEMATICS AND SOFTWARE ENGINEERING, UNIVERSITY OF DETROIT
MERCY, 4001 WEST McNICHOLS ROAD, DETROIT MI 48221-3038, USA
E-mail address: `archeyde@udmercy.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OREGON, EUGENE OR 97403-1222,
USA